

Rabbit HTT Polyclonal Antibody | anti-HTT antibody

HTT Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA314621	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HTT; Polyclonal Antibody; HTT Rabbit Polyclonal Antibody; HD; IT15; Huntingtin; anti-HTT antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applicable Applications for anti-HTT antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human HTT.
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.


IMAGES


IF (Immunofluorescence)

Immunofluorescence analysis of U-251 MG cells using HTT Rabbit pAb.


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat brain using HTT Rabbit pAb.



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using HTT antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-HTT antibody	Huntingtin is a disease gene linked to Huntington's disease, a neurodegenerative disorder characterized by loss of striatal neurons. This is thought to be caused by an expanded, unstable trinucleotide repeat in the huntingtin gene, which translates as a polyglutamine repeat in the protein product. A fairly broad range of trinucleotide repeats (9-35) has been identified in normal controls, and repeat numbers in excess of 40 have been described as pathological. The huntingtin locus is large, spanning 180 kb and consisting of 67 exons. The huntingtin gene is widely expressed and is required for normal development. It is expressed as 2 alternatively polyadenylated forms displaying different relative abundance in various fetal and adult tissues. The larger transcript is approximately 13.7 kb and is expressed predominantly in adult and fetal brain whereas the smaller transcript of approximately 10.3 kb is more widely expressed. The genetic defect leading to Huntington's disease may not necessarily eliminate transcription, but may confer a new property on the mRNA or alter the function of the protein. One candidate is the huntingtin-associated protein-1, highly expressed in brain, which has increased affinity for huntingtin protein with expanded polyglutamine repeats. This gene contains an upstream open reading frame in the 5' UTR that inhibits expression of the huntingtin gene product through translational repression.
Product Categories/Family for anti-HTT antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	90903231
NCBI GeneID	3064
NCBI Accession #	NP_002102.4
NCBI GenBank Nucleotide #	NM_002111.7
UniProt Accession #	P42858 ; Q9UQB7
Molecular Weight	347,603 Da
NCBI Official Full Name	huntingtin
NCBI Official Synonym Full Names	huntingtin
NCBI Official Symbol	HTT
NCBI Official Synonym Symbols	HD; IT15
NCBI Protein Information	huntingtin; huntington disease protein
UniProt Protein Name	Huntingtin
UniProt Gene Name	HTT
UniProt Synonym Gene Names	HD; IT15; HD protein

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/htt-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/314621>