

HDAC2 recombinant antibody

HDAC2 Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA315014	Rabbit	Human, Mouse, Rat	Flow Cytometry, Functional Assay, Immunoprecipitation, Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HDAC2; Monoclonal Recombinant Antibody; HDAC2 Rabbit mAb; HD2; Histone deacetylase 2; RPD3; YAF1; YY1 associated factor 1; HDAC2 recombinant antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	Rabbit IgG
Specificity	HDAC2 Antibody detects endogenous levels of total HDAC2
Purity/Purification	Affinity-chromatography
Form/Format	Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Applicable Applications for HDAC2 recombinant antibody	FCM/FACS (Flow Cytometry), IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human HDAC2
Preparation and Storage	Store at +4 degree C short term. Store at -20 degree C long term. Avoid freeze/thaw cycle.

IMAGES


IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human breast, using HDAC2 Antibody.


WB (Western Blot)

Western blot analysis of HDAC2 expression in K562 cell lysate.

ADDITIONAL INFORMATION

Related Product Information for HDAC2 recombinant antibody	In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino-terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA.
Product Categories/Family for HDAC2 recombinant antibody	Recombinant Rabbit Monoclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	293336691
NCBI GeneID	3066
NCBI Accession #	NP_001518.3
NCBI GenBank Nucleotide #	NM_001527.3
UniProt Accession #	Q92769; Q5SRI8; Q5SZ86; Q8NEH4; B4DL58; E1P561
Molecular Weight	55,364 Da
NCBI Official Full Name	histone deacetylase 2
NCBI Official Synonym Full Names	histone deacetylase 2
NCBI Official Symbol	HDAC2
NCBI Official Synonym Symbols	HD2; RPD3; YAF1
NCBI Protein Information	histone deacetylase 2; YY1-associated factor 1; transcriptional regulator homolog RPD3
UniProt Protein Name	Histone deacetylase 2
UniProt Gene Name	HDAC2
UniProt Synonym Gene Names	HD2
UniProt Entry Name	HDAC2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/hdac2-recombinant-antibody-flow-cytometry-immunoprecipitation-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/315014>