

Rabbit CD274 Polyclonal Antibody | anti-CD274 antibody

CD274 Rabbit Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA318806	Rabbit	Human, Rat, Mouse	ELISA, Immunofluorescence, Immunohistochemistry


PRODUCT SYNONYMS

CD274; Polyclonal Antibody; CD274 Rabbit Polyclonal Antibody; Programmed cell death 1 ligand 1 (PD-L1)(PDCD1 ligand 1)(Programmed death ligand 1)(B7 homolog 1)(B7-H1)(CD antigen CD274); BH6842; anti-CD274 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Specificity	The antibody detects endogenous CD274
Purity/Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-CD274 antibody	ELISA, IF (Immunofluorescence), IHC (Immunohistochemistry)
Immunogen	Synthetic peptide from human protein at AA range: 181-230
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-CD274 antibody	This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	390979639
NCBI GeneID	29126
NCBI Accession #	NP_001254635.1
NCBI GenBank Nucleotide #	NM_001267706.1
UniProt Accession #	Q9NZQ7; Q14CJ2; Q2V8D5; Q66RK1; Q6WEX4; Q9NUZ5; B2RBA2; B4DU27
Molecular Weight	33,275 Da
NCBI Official Full Name	programmed cell death 1 ligand 1 isoform b
NCBI Official Synonym Full Names	CD274 molecule
NCBI Official Symbol	CD274
NCBI Official Synonym Symbols	B7-H; B7H1; PDL1; PD-L1; PDCD1L1; PDCD1LG1
NCBI Protein Information	programmed cell death 1 ligand 1; B7 homolog 1; CD274 antigen; PDCD1 ligand 1; programmed death ligand 1
UniProt Protein Name	Programmed cell death 1 ligand 1
UniProt Gene Name	CD274
UniProt Synonym Gene Names	B7H1; PDCD1L1; PDCD1LG1; PDL1; PD-L1; PDCD1 ligand 1; Programmed death ligand 1; B7-H1
UniProt Entry Name	PD1L1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cd274-antibody-immunofluorescence-immunohistochemistry-elisa/318806>