

Mouse anti-Rabbit IgG Monoclonal Antibody | anti-IgG antibody

Rabbit IgG Mouse mAb

CATALOG # AAA320394	HOST Mouse	REACTIVITY Rabbit	APPLICATIONS ELISA, Western Blot
-------------------------------	----------------------	-----------------------------	--


PRODUCT SYNONYMS

IgG; Monoclonal Antibody; Rabbit IgG Mouse mAb; Antibody; Ig gamma chain C region; Ig kappa-b4 chain C region; Ig kappa-b5 chain C region; Ig kappa-b9 chain C region; Ig lambda chain C region; K-BAS.; anti-IgG antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Rabbit
Clonality	Monoclonal
Specificity	This antibody detects rabbit IgG.
Purity/Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-IgG antibody	ELISA, WB (Western Blot)
Immunogen	Rabbit IgG
Modification	Unmodified/Total
Preparation and Storage	Store at -20 degree C/1 year

IMAGES




WB (Western Blot)

Western blot detection of MEK-2 in THP-1, Jurkat, Raji, 3T3 and Hela cell lysates using 310001 (MEK-2 rabbit pAb, 1:1000 diluted), then incubated with 201069 (Rabbit IgG mAb, 1:50000 diluted). Anti-mouse IgG, HRP-linked Antibody was used for detecting. Predicted band size: 44 kDa. Observed band size: 44 kDa.



WB (Western Blot)

Western blot analysis of decreasing concentrations of total rabbit IgG reduced and denatured in 2X SDS loading buffer with DTT using Mouse Anti-Rabbit IgG (Heavy-Chain Specific) mAb (1:1000 diluted).



IP (Immunoprecipitation)

Immunoprecipitation analysis of CHO-K1 cell lysates transfected with HA-tagged NGAL using HA-Tag Rabbit pAb (390001.1:100 diluted)(lane 2). Lane 1 contains rabbit IgG IP control lysate. All lanes are probed with HA-Tag Rabbit pAb (390001.1:1000 diluted) as the primary antibody. Panel A incubated with secondary antibodies Anti-Rabbit whole IgG. HRP-linked Antibody for detecting. Panel B incubated with Anti-Rabbit IgG (Heavy-Chain Specific) mAb(1:50000 diluted) and the Anti-mouse whole IgG. HRP-linked Antibody was used for detecting.

ADDITIONAL INFORMATION

Product Categories/Family for anti-IgG antibody	Primary Antibodies
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.	

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/monoclonal/igg-antibody-western-blot-elisa/320394>