

Rabbit RSK2 Polyclonal Antibody | anti-RSK2 antibody

Phospho-RSK2 (Ser227) Antibody


CATALOG # AAA321383	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---

PRODUCT SYNONYMS


RSK2; Polyclonal Antibody; Phospho-RSK2 (Ser227) Antibody; 90 kDa ribosomal protein S6 kinase 3; CLS; HU 3; HU2; HU3; Insulin stimulated protein kinase 1; Insulin-stimulated protein kinase 1; ISPK-1; ISPK1; KS6A3_HUMAN; MAP kinase activated protein kinase 1b; MAP kinase-activated protein kinase 1b; MAPK activated protein kinase 1b; MAPK-activated protein kinase 1b; MAPKAP kinase 1b; MAPKAPK 1b; MAPKAPK-1b; MAPKAPK1B; Mental retardation; X linked 19; MRX19; OTTHUMP00000023036; p90 RSK2; p90 RSK3; p90-RSK 3; p90RSK3; pp90RSK2; Ribosomal protein S6 kinase 90kDa polypeptide 3; Ribosomal protein S6 kinase alpha 3; Ribosomal protein S6 kinase alpha-3; Ribosomal protein s6 kinase ii alpha 2; Ribosomal S6 kinase 2; Rps6ka3; RSK; RSK-2; S6 kinase 2; S6K alpha3; S6K-alpha-3; anti-RSK2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Phospho-RSK2 (Ser227) antibody detects endogenous levels of RSK2.
Purity/Purification	From purified rabbit serum by affinity purification via sequential chromatography on phospho-and non-phospho-peptide affinity columns.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	740
Applicable Applications for anti-RSK2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human RSK2 around the phosphorylation site of Ser227.
Subcellular Location	Nucleus
Tissue Specificity	Expressed in many tissues, highest levels in skeletal muscle.
Predicted Cross Reactivity	Pig, Bovine, Horse, Sheep, Rabbit, Dog, Chicken, Xenopus
Similarity	Pig (100%), Bovine (100%), Horse (100%), Sheep (100%), Rabbit (100%), Dog (100%), Chicken (100%), Xenopus (100%)
Conjugation	Unconjugated
Epitope	Phospho Ser227
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



WB (Western Blot)
Western blot analysis of extracts of hela cells treated with TPA/UV, using Phospho-RSK2 (Ser227) Antibody.



IHC (Immunohistochemistry)
AAA321383 at 1/100 staining Mouse heart tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

ADDITIONAL INFORMATION

Related Product Information for anti-RSK2 antibody	Function: Serine/threonine-protein kinase that acts downstream of ERK (MAPK1/ERK2 and MAPK3/ERK1) signaling and mediates mitogenic and stress-induced activation of the transcription factors CREB1, ETV1/ER81 and NR4A1/NUR77, regulates translation through RPS6 and EIF4B phosphorylation, and mediates cellular proliferation, survival, and differentiation by modulating mTOR signaling and repressing pro-apoptotic function of BAD and DAPK1. In fibroblast, is required for EGF-stimulated phosphorylation of CREB1 and histone H3 at 'Ser-10', which results in the subsequent transcriptional activation of several immediate-early genes. In response to mitogenic stimulation (EGF and PMA), phosphorylates and activates NR4A1/NUR77 and ETV1/ER81 transcription factors and the cofactor CREBBP. Upon insulin-derived signal, acts indirectly on the transcription regulation of several genes by phosphorylating GSK3B at 'Ser-9' and inhibiting its activity. Phosphorylates RPS6 in response to serum or EGF via an mTOR-independent mechanism and promotes translation initiation by facilitating assembly of the preinitiation complex. In response to insulin, phosphorylates EIF4B, enhancing EIF4B affinity for the EIF3 complex and stimulating cap-dependent translation. Is involved in the mTOR nutrient-sensing pathway by directly phosphorylating TSC2 at 'Ser-1798', which potently inhibits TSC2 ability to suppress mTOR signaling, and mediates phosphorylation of RPTOR, which regulates mTORC1 activity and may promote rapamycin-sensitive signaling independently of the PI3K/AKT pathway. Mediates cell survival by phosphorylating the pro-apoptotic proteins BAD and DAPK1 and suppressing their pro-apoptotic function. Promotes the survival of hepatic stellate cells by phosphorylating CEBPB in response to the hepatotoxin carbon tetrachloride (CCl4). Is involved in cell cycle regulation by phosphorylating the CDK inhibitor CDKN1B, which promotes CDKN1B association with 14-3-3 proteins and prevents its translocation to the nucleus and inhibition of G1 progression. In LPS-stimulated dendritic cells, is involved in TLR4-induced macropinocytosis, and in myeloma cells, acts as effector of FGFR3-mediated transformation signaling, after direct phosphorylation at Tyr-529 by FGFR3. Negatively regulates EGF-induced MAPK1/3 phosphorylation via phosphorylation of SOS1. Phosphorylates SOS1 at 'Ser-1134' and 'Ser-1161' that create YWHAB and YWHAE binding sites and which contribute to the negative regulation of MAPK1/3 phosphorylation (By similarity). Phosphorylates EPHA2 at 'Ser-897', the RPS6KA-EPHA2 signaling pathway controls cell migration (PubMed:26158630). Subunit Structure: Forms a complex with either MAPK1/ERK2 or MAPK3/ERK1 in quiescent cells. Transiently dissociates following mitogenic stimulation (By similarity). Interacts with NFATC4, ETV1/ER81 and FGFR1. Post-translational Modifications: Activated by phosphorylation at Ser-227 by PDPK1. Autophosphorylated on Ser-386, as part of the activation process. May be phosphorylated at Thr-365 and Ser-369 by MAPK1/ERK2 and MAPK3/ERK1. Can also be activated via phosphorylation at Ser-386 by MAPKAPK2. N-terminal myristoylation results in an activated kinase in the absence of added growth factors. Similarity: Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4759050
NCBI GeneID	6197
NCBI Accession #	NP_004577.1

NCBI GenBank Nucleotide #	NM_004586.2
UniProt Accession #	P51812 ; Q4VAP3 ; Q59H26 ; Q5JPK8 ; Q7Z3Z7 ; B2R9V4
Molecular Weight	Observed: 90 kDa Predicted: 84 kDa
NCBI Official Full Name	ribosomal protein S6 kinase alpha-3
NCBI Official Synonym Full Names	ribosomal protein S6 kinase A3
NCBI Official Symbol	RPS6KA3
NCBI Official Synonym Symbols	CLS; RSK; HU-3; RSK2; MRX19; ISPK-1; p90-RSK2; pp90RSK2; MAPKAPK1B; S6K-alpha3
NCBI Protein Information	ribosomal protein S6 kinase alpha-3
UniProt Protein Name	Ribosomal protein S6 kinase alpha-3
UniProt Gene Name	RPS6KA3
UniProt Synonym Gene Names	ISPK1; MAPKAPK1B; RSK2; S6K-alpha-3; p90-RSK 3; p90RSK3; ISPK-1; MAPK-activated protein kinase 1b; MAPKAP kinase 1b; MAPKAPK-1b; RSK-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/rsk2-antibody-immunohistochemistry-western-blot-elisa/321383>