

Rabbit anti-Human, Mouse E2F2 Polyclonal Antibody | anti-E2F2 antibody

E2F2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA323975	Rabbit	Human, Mouse	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

E2F2; Polyclonal Antibody; E2F2 Antibody; dE2F2; E2F transcription factor 2; E2F-2; E2F2_HUMAN; Transcription factor E2F2; anti-E2F2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	E2F2 antibody detects endogenous levels of total E2F2
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	437
Applicable Applications for anti-E2F2 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Nucleus.
Tissue Specificity	Highest level of expression is found in placenta, low levels are found in lung. Found as well in many immortalized cell lines derived from tumor samples.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)
Western blot analysis of extracts from HepG2 cells using E2F2 antibody.



IF (Immunofluorescence)
AAA323975 staining HepG2 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody, diluted at 1/600, was used as secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-E2F2 antibody	Function: Transcription activator that binds DNA cooperatively with DP proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from g1 to s phase. E2F2 binds specifically to RB1 in a cell-cycle dependent manner. Subunit Structure: Component of the DRTF1/E2F transcription factor complex. Forms heterodimers with DP family members. The E2F2 complex binds specifically hypophosphorylated retinoblastoma protein RB1. During the cell cycle, RB1 becomes phosphorylated in mid-to-late G1 phase, detaches from the DRTF1/E2F complex, rendering E2F transcriptionally active. Viral oncoproteins, notably E1A, T-antigen and HPV E7, are capable of sequestering RB1, thus releasing the active complex. Binds EAPP. Post-translational Modifications: Phosphorylated by CDK2 and cyclin A-CDK2 in the S-phase. Similarity: Belongs to the E2F/DP family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758226
NCBI GeneID	1870
NCBI Accession #	NP_004082.1
NCBI GenBank Nucleotide #	NM_004091.3
UniProt Accession #	Q14209; Q7Z6H1; B2R9W1
Molecular Weight	Observed: 48 kD Predicted: 48 kDa
NCBI Official Full Name	transcription factor E2F2
NCBI Official Synonym Full Names	E2F transcription factor 2
NCBI Official Symbol	E2F2
NCBI Official Synonym Symbols	E2F-2
NCBI Protein Information	transcription factor E2F2
UniProt Protein Name	Transcription factor E2F2
UniProt Gene Name	E2F2
UniProt Synonym Gene Names	E2F-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/e2f2-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/323975>