

Rabbit P2RY4 Polyclonal Antibody | anti-P2RY4 antibody

P2RY4 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA325739	Rabbit	Human, Mouse, Rat	ELISA, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

P2RY4; Polyclonal Antibody; P2RY4 Antibody; NRU; Nucleotide receptor uridine; P2P; P2ry4; P2RY4_HUMAN; P2Y purinoceptor 4; P2Y4; P2Y4R; Pyrimidinergic receptor P2Y G protein coupled 4; UNR; Uridine nucleotide receptor; anti-P2RY4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	P2RY4 antibody detects endogenous levels of total P2RY4
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	365
Applicable Applications for anti-P2RY4 antibody	ELISA, ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	A synthesized peptide
Subcellular Location	Cell Membrane
Tissue Specificity	Pancreas.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)
Western blot analysis of extracts from HeLa cells, using P2RY4 antibody.



IF (Immunofluorescence)
AAA325739 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25 degree C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37 degree C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-P2RY4 antibody	Function: Receptor for UTP and UDP coupled to G-proteins that activate a phosphatidylinositol-calcium second messenger system. Not activated by ATP or ADP. Post-translational Modifications: Phosphorylation of Ser-333 and Ser-334 is a key step in agonist-dependent desensitization and loss of surface P2RY4. This phosphorylation does not involve PKC, nor other calcium activated kinases. Similarity: Belongs to the G-protein coupled receptor 1 family.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505561
NCBI GeneID	5030
NCBI Accession #	NP_002556.1
NCBI GenBank Nucleotide #	NM_002565.3
UniProt Accession #	P51582; Q4VBB7; Q4VBB8; Q502W2; Q5JT22
Molecular Weight	Observed: 41 kD Predicted: 41 kDa
NCBI Official Full Name	P2Y purinoceptor 4
NCBI Official Synonym Full Names	pyrimidinergic receptor P2Y4
NCBI Official Symbol	P2RY4
NCBI Official Synonym Symbols	NRU; P2P; UNR; P2Y4
NCBI Protein Information	P2Y purinoceptor 4
UniProt Protein Name	P2Y purinoceptor 4
UniProt Gene Name	P2RY4
UniProt Synonym Gene Names	NRU; P2Y4; UNR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.