

Rabbit PTHLH Polyclonal Antibody | anti-PTHLH antibody

PTHLH Antibody



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA326298	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

PTHLH; Polyclonal Antibody; PTHLH Antibody; HHM; Humoral hypercalcemia of malignancy; MGC14611; Osteostatin; Parathyroid hormone like hormone; Parathyroid hormone like protein; Parathyroid hormone like related protein; Parathyroid like protein; PLP; PTH related protein; PTHR; PTHRP; anti-PTHLH antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	PTHLH antibody detects endogenous levels of total PTHLH
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	175
Applicable Applications for anti-PTHLH antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human PTHLH
Subcellular Location	Cytoplasm. Nucleus. Secreted.
Tissue Specificity	Ubiquitous. Also expressed in the mammary gland.
Predicted Cross Reactivity	Bovine, Horse, Sheep, Rabbit, Dog, Chicken
Similarity	Bovine (92%), Horse (100%), Sheep (92%), Rabbit (100%), Dog (100%), Chicken (92%)
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



IHC (Immunohistochemistry)
AAA326298 at 1/100 staining Mouse colon tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

WB (Western Blot)
Western blot analysis of extracts of A549, using PTHLH antibody. The lane on the left is treated with the antigen-specific peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-PTHLH antibody

Description: The protein encoded by this gene is a member of the parathyroid hormone family. This hormone regulates endochondral bone development and epithelial-mesenchymal interactions during the formation of the mammary glands and teeth. This hormone is involved in lactation possibly by regulating the mobilization and transfer of calcium to the milk. The receptor of this hormone, PTHR1, is responsible for most cases of humoral hypercalcemia of malignancy. Four alternatively spliced transcript variants encoding two distinct isoforms have been observed. There is also evidence for alternative translation initiation from non-AUG (CUG and GUG) start sites, in-frame and downstream of the initiator AUG codon, to give rise to nuclear forms of this hormone. [provided by RefSeq, Jul 2008]

Function: Neuroendocrine peptide which is a critical regulator of cellular and organ growth, development, migration, differentiation and survival and of epithelial calcium ion transport. Regulates endochondral bone development and epithelial-mesenchymal interactions during the formation of the mammary glands and teeth. Required for skeletal homeostasis. Promotes mammary mesenchyme differentiation and bud outgrowth by modulating mesenchymal cell responsiveness to BMPs. Upregulates BMPR1A expression in the mammary mesenchyme and this increases the sensitivity of these cells to BMPs and allows them to respond to BMP4 in a paracrine and/or autocrine fashion. BMP4 signaling in the mesenchyme, in turn, triggers epithelial outgrowth and augments MSX2 expression, which causes the mammary mesenchyme to inhibit hair follicle formation within the nipple sheath (By similarity). Promotes colon cancer cell migration and invasion in an integrin alpha-6/beta-1-dependent manner through activation of Rac1.

Subunit Structure: PTHrP interacts with PTH1R (via N-terminal extracellular domain).

Post-translational Modifications: There are 3 principal secretory forms, called PTHrP[1-36], PTHrP[38-94], and osteostatin (PTHrP[107-139]) arising from endoproteolytic cleavage of the initial translation product. Each of these secretory forms is believed to have one or more of its own receptors that mediates the normal paracrine, autocrine and endocrine actions.

Similarity: Belongs to the parathyroid hormone family.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506269
NCBI GeneID	5744
NCBI Accession #	NP_002811.1
NCBI GenBank Nucleotide #	NM_002820.2
UniProt Accession #	P12272; Q15251; Q6FH74
Molecular Weight	Observed: 20 kDa Predicted: 21 kDa
NCBI Official Full Name	parathyroid hormone-related protein isoform 2 preproprotein
NCBI Official Synonym Full Names	parathyroid hormone like hormone

NCBI Official Symbol	PTHLH
NCBI Official Synonym Symbols	HHM; PLP; BDE2; PTHR; PTHRP
NCBI Protein Information	parathyroid hormone-related protein
UniProt Protein Name	Parathyroid hormone-related protein
UniProt Gene Name	PTHLH
UniProt Synonym Gene Names	PTHRP; PTH-rP; PTHrP; PLP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/ptlh-antibody-immunohistochemistry-western-blot-elisa/326298>