

Rabbit USP7 Polyclonal Antibody | anti-USP7 antibody

USP7 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA327199	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS


USP7; Polyclonal Antibody; USP7 Antibody; Deubiquitinating enzyme 7; HAUSP; Herpes virus associated ubiquitin specific protease; Herpesvirus-associated ubiquitin-specific protease; TEF 1; tef-1; TEF1; Ubiquitin carboxyl terminal hydrolase 7; Ubiquitin carboxyl-terminal hydrolase 7; Ubiquitin specific peptidase 7 (herpes virus associated); Ubiquitin specific peptidase 7; Ubiquitin specific peptidase 7 herpes virus associated; Ubiquitin specific processing protease 7; Ubiquitin specific protease 7 (herpes virus associated); Ubiquitin specific protease 7; Ubiquitin specific protease 7 herpes virus associated; Ubiquitin thioesterase 7; Ubiquitin thiolesterase 7; Ubiquitin-specific-processing protease 7; UBP 7; UBP-7; UBP7; UBP7_HUMAN; USP 7; usp-7; Usp7; VMW110-ASSOCIATED PROTEIN; 135-KD; anti-USP7 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	USP7 antibody detects endogenous levels of total USP7
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	1044
Applicable Applications for anti-USP7 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthesized peptide derived from human USP7
Subcellular Location	Nucleus. Cytoplasm. Nucleus, PML Body. Present in a minority of ND10 nuclear bodies. Association with ICP0/VMW110 at early times of infection leads to an increased proportion of USP7-containing ND10. Colocalizes with ATXN1 in the nucleus. Colocalized with DAXX in speckled structures. Colocalized with PML and PTEN in promyelocytic leukemia protein (PML) nuclear bodies.
Tissue Specificity	Widely expressed. Overexpressed in prostate cancer.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.



IHC (Immunohistochemistry)
AAA327199 at 1/100 staining Human breast cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22 degree C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



WB (Western Blot)
Western blot analysis USP7 using COLO205 whole cell lysates

ADDITIONAL INFORMATION

Related Product Information for anti-USP7 antibody	Function: Hydrolase that deubiquitinates target proteins such as FOXO4, p53/TP53, MDM2, ERCC6, DNMT1, UHRF1, PTEN, KMT2E/MLL5 and DAXX (PubMed:11923872, PubMed:15053880, PubMed:16964248, PubMed:18716620, PubMed:25283148, PubMed:26678539). Together with DAXX, prevents MDM2 self-ubiquitination and enhances the E3 ligase activity of MDM2 towards p53/TP53, thereby promoting p53/TP53 ubiquitination and proteasomal degradation (PubMed:15053880, PubMed:16845383, PubMed:18566590, PubMed:20153724). Deubiquitinates p53/TP53, preventing degradation of p53/TP53, and enhances p53/TP53-dependent transcription regulation, cell growth repression and apoptosis (PubMed:25283148). Deubiquitinates p53/TP53 and MDM2 and strongly stabilizes p53/TP53 even in the presence of excess MDM2, and also induces p53/TP53-dependent cell growth repression and apoptosis (PubMed:11923872). Deubiquitination of FOXO4 in presence of hydrogen peroxide is not dependent on p53/TP53 and inhibits FOXO4-induced transcriptional activity (PubMed:16964248). In association with DAXX, is involved in the deubiquitination and translocation of PTEN from the nucleus to the cytoplasm, both processes that are counteracted by PML (PubMed:18716620). Deubiquitinates KMT2E/MLL5 preventing KMT2E/MLL5 proteasomal-mediated degradation (PubMed:26678539). Involved in cell proliferation during early embryonic development. Involved in transcription-coupled nucleotide excision repair (TC-NER) in response to UV damage: recruited to DNA damage sites following interaction with KIAA1530/UVSSA and promotes deubiquitination of ERCC6, preventing UV-induced degradation of ERCC6 (PubMed:22466611, PubMed:22466612). Involved in maintenance of DNA methylation via its interaction with UHRF1 and DNMT1: acts by mediating deubiquitination of UHRF1 and DNMT1, preventing their degradation and promoting DNA methylation by DNMT1 (PubMed:21745816, PubMed:22411829). Deubiquitinates alkylation repair enzyme ALKBH3. OTUD4 recruits USP7 and USP9X to stabilize ALKBH3, thereby promoting the repair of alkylated DNA lesions (PubMed:25944111). Acts as a chromatin regulator via its association with the Polycomb group (PcG) multiprotein PRC1-like complex; may act by deubiquitinating components of the PRC1-like complex (PubMed:20601937). Able to mediate deubiquitination of histone H2B; it is however unsure whether this activity takes place in vivo (PubMed:20601937). Exhibits a preference towards 'Lys-48'-linked ubiquitin chains (PubMed:22689415). Increases regulatory T-cells (Treg) suppressive capacity by deubiquitinating and stabilizing the transcription factor FOXP3 which is crucial for Treg cell function (PubMed:23973222). Subunit Structure: Monomer. Homodimer. Part of a complex with DAXX, MDM2, RASSF1 and USP7 (PubMed:18566590). Part of a complex with DAXX, MDM2 and USP7 (PubMed:16845383). Interacts with MDM2; the interaction is independent of p53/TP53. Interacts with DAXX; the interaction is direct and independent of MDM2 and p53/TP53 (PubMed:16845383). Component of a complex composed of KMT2E/MLL5 (isoform 3), OGT (isoform 1) and USP7; the complex stabilizes KMT2E/MLL5, preventing KMT2E/MLL5 ubiquitination and proteasomal-mediated degradation (PubMed:26678539). Interacts (via MATH domain) with KMT2E/MLL5 isoform 3 (PubMed:26678539). Interacts with OGT isoform 1 (PubMed:26678539). Interacts with FOXO4; the interaction is enhanced in presence of hydrogen peroxide and occurs independently of p53/TP53 (PubMed:16964248). Interacts with p53/TP53; the interaction is enhanced in response to DNA damage (PubMed:25283148). Interacts with TSPYL5; this impairs interaction with p53/TP53 (PubMed:21170034). Interacts with PTEN; the interaction is direct (PubMed:18716620). Interacts with ATXN1 and the strength of interaction is influenced by the length of the poly-Gln region in ATXN1 (PubMed:12093161). A weaker interaction seen with mutants having longer poly-Gln regions (PubMed:12093161). Interacts with KIAA1530/UVSSA (PubMed:22466611, PubMed:22466612). Interacts with ABRAXAS2; the interaction is direct (PubMed:25283148). Identified in a complex with TP53/p53 and ABRAXAS2 (PubMed:25283148). Interacts with MEX3C and antagonizes its ability to degrade mRNA (PubMed:22863774). Interacts with DNMT1 and UHRF1 (PubMed:21745816, PubMed:22411829). Interacts with FOXP3 (PubMed:23973222). Interacts (via MATH domain) with RNF220. Associated component of the Polycomb group (PcG) multiprotein PRC1-like complex (PubMed:20601937). Interacts with EPOP (By similarity). Interacts with OTUD4 and USP9X; the interaction is direct (PubMed:25944111). Post-translational Modifications: Isoform 1: Phosphorylated. Isoform 1 is phosphorylated at positions Ser-18 and Ser-963. Isoform 2: Not phosphorylated. Isoform 1: Polyubiquitinated. Isoform 2: Not Polyubiquitinated. Isoform 1 and isoform 2: Not sumoylated. Isoform 1 and isoform 2: Polyubiquitinated by herpesvirus 1 trans-acting transcriptional protein ICP0/VMW110; leading to its subsequent proteasomal degradation. Isoform 1: Ubiquitinated at Lys-869. Similarity: The C-terminus plays a role in its oligomerization. Belongs to the peptidase C19 family.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	1013061510
NCBI GeneID	7874
NCBI Accession #	NP_001308787.1
NCBI GenBank Nucleotide #	NM_001321858.1
UniProt Accession #	Q93009; A6NMY8; B7Z815; H0Y3G8

Molecular Weight	Observed: 128 kDa Predicted: 129 kDa
NCBI Official Full Name	ubiquitin carboxyl-terminal hydrolase 7 isoform 4
NCBI Official Synonym Full Names	ubiquitin specific peptidase 7
NCBI Official Symbol	USP7
NCBI Official Synonym Symbols	TEF1; HAUSP
NCBI Protein Information	ubiquitin carboxyl-terminal hydrolase 7
UniProt Protein Name	Ubiquitin carboxyl-terminal hydrolase 7
UniProt Gene Name	USP7

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/usp7-antibody-immunohistochemistry-western-blot-elisa/327199>