

Rabbit Cofilin Polyclonal Antibody | anti-CFL1 antibody

Cofilin Antibody

CATALOG # AAA327569	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA, Western Blot
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

Cofilin; Polyclonal Antibody; Cofilin Antibody; 18 kDa phosphoprotein; CFL 1; CFL; CFL1; COF1_HUMAN; Cofilin 1; Cofilin 1 non muscle; Cofilin non muscle isoform; Cofilin-1; epididymis secretory protein Li 15; HEL-S-15; non-muscle isoform; p18; CFL 2; CFL2; COF2_HUMAN; Cofilin 2 muscle; Cofilin muscle; Cofilin muscle isoform; Cofilin-2; Cofilin2; muscle isoform; NEM 7; NEM7; anti-CFL1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	Cofilin antibody detects endogenous levels of total Cofilin
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin.
Form/Format	Liquid Phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1mg/ml (varies by lot)
Sequence Length	166
Applicable Applications for anti-CFL1 antibody	ELISA, WB (Western Blot)
Immunogen	A synthesized peptide derived from human Cofilin
Subcellular Location	Nucleus Matrix. Cytoplasm > Cytoskeleton. Almost completely in nucleus in cells exposed to heat shock or 10% dimethyl sulfoxide.
Tissue Specificity	Widely distributed in various tissues.
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.

IMAGES



WB (Western Blot)
Western blot analysis of Cofilin using A549 whole cell lysates



WB (Western Blot)
Western blot analysis of extracts of various tissue sample, using Cofilin Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CFL1 antibody	Function: Binds to F-actin and exhibits pH-sensitive F-actin depolymerizing activity. Regulates actin cytoskeleton dynamics. Important for normal progress through mitosis and normal cytokinesis. Plays a role in the regulation of cell morphology and cytoskeletal organization. Required for the up-regulation of atypical chemokine receptor ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation (PubMed:11812157, PubMed:15580268, PubMed:21834987, PubMed:23633677). Required for neural tube morphogenesis and neural crest cell migration (By similarity). Subunit Structure: Can bind G-and F-actin in a 1:1 ratio of cofilin to actin. It is a major component of intranuclear and cytoplasmic actin rods. Post-translational Modifications: Inactivated by phosphorylation on Ser-3. Phosphorylated on Ser-3 in resting cells (By similarity). Dephosphorylated by PDXP/chronophin; this restores its activity in promoting actin filament depolymerization. The phosphorylation of Ser-24 may prevent recognition of the nuclear localization signal (By similarity). Phosphorylated via a ARRB1-RAC1-LIMK1-PAK1 cascade upon active ligand stimulation of atypical chemokine receptor ACKR2. Similarity: Belongs to the actin-binding proteins ADF family.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	5031635
NCBI GeneID	1072
NCBI Accession #	NP_005498.1
NCBI GenBank Nucleotide #	NM_005507.2
UniProt Accession #	P23528; Q53Y87; Q9UCA2; B3KUQ1
Molecular Weight	Observed: 18 kDa Predicted: 19 kDa
NCBI Official Full Name	cofilin-1
NCBI Official Synonym Full Names	cofilin 1
NCBI Official Symbol	CFL1
NCBI Official Synonym Symbols	CFL; cofilin; HEL-S-15
NCBI Protein Information	cofilin-1
UniProt Protein Name	Cofilin-1
UniProt Gene Name	CFL1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cofilin-antibody-western-blot-elisa/327569>