

Rabbit Apolipoprotein M Polyclonal Antibody | anti-APOM antibody

Apolipoprotein M Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA329145	Rabbit	Human, Mouse, Rat Predicted Reactivity: Pig(100%), Bovine(100%), Horse(100%), Sheep(100%), Dog(88%)	ELISA


PRODUCT SYNONYMS

Apolipoprotein M; Polyclonal Antibody; Apolipoprotein M Antibody; Apo M; Apo-M; ApoM; APOM_HUMAN; G3A; HSPC336; MGC22400; NG20; NG20 like protein; Protein G3a; anti-APOM antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat Predicted Reactivity: Pig(100%), Bovine(100%), Horse(100%), Sheep(100%), Dog(88%)
Clonality	Polyclonal
Isotype	IgG
Specificity	Apolipoprotein M Antibody detects endogenous levels of total Apolipoprotein M.
Purity/Purification	The antiserum was purified by peptide affinity chromatography using SulfoLink Coupling Resin (Thermo Fisher Scientific).
Form/Format	Liquid. Rabbit IgG in phosphate buffered saline, pH7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol
Concentration	1mg/ml (varies by lot)
Sequence Length	116
Applicable Applications for anti-APOM antibody	ELISA
Immunogen	A synthesized peptide derived from human Apolipoprotein M, corresponding to a region within the internal amino acids.
Fragment	Fab fragment
Preparation and Storage	Store at -20 degree C. Stable for 12 months from date of receipt.


IMAGES



IF (Immunofluorescence)
AAA329145 staining HeLa cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(AAA329145 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary antibody. The nuclear counter stain is DAPI(blue).



WB (Western Blot)
Western blot analysis of extracts from Mouse heart, using Apolipoprotein M Antibody. The lane on the left was treated with blocking peptide.



WB (Western Blot)
Western blot analysis of extracts from Human Plasma, using Apolipoprotein M Antibody. The lane on the left was treated with blocking peptide.

ADDITIONAL INFORMATION

Related Product Information for anti-APOM antibody	Function: Probably involved in lipid transport. Can bind sphingosine-1-phosphate, myristic acid, palmitic acid and stearic acid, retinol, all-trans-retinoic acid and 9-cis-retinoic acid.
	Subcellular Location: Secreted. Note: Present in high density lipoprotein (HDL) and to a lesser extent in triglyceride-rich lipoproteins (TGRLP) and low density lipoproteins (LDL).
	Tissue Specificity: Plasma protein. Expressed in liver and kidney.
	Subunit Structure: Interacts with LRP2; LRP2 mediates APOM renal uptake and subsequent lysosomal degradation.
	Similarity: Belongs to the calycin superfamily. Lipocalin family. Highly divergent.

NCBI AND UNIPROT INFORMATION

NCBI GI #	371873523
NCBI GeneID	55937
NCBI Accession #	NP_001243098.1

NCBI GenBank Nucleotide #	NP_001243098.1
UniProt Accession #	O95445
Molecular Weight	Observed Molecular Weight: (Observed)21kD. Predicted Molecular Weight: (Calculated)21kDa.
NCBI Official Full Name	apolipoprotein M isoform 2
NCBI Official Synonym Full Names	apolipoprotein M
NCBI Official Symbol	APOM
NCBI Official Synonym Symbols	G3a; NG20; apo-M; HSPC336
NCBI Protein Information	apolipoprotein M
UniProt Protein Name	Apolipoprotein M
UniProt Gene Name	APOM
UniProt Synonym Gene Names	G3A; NG20; Apo-M; ApoM
UniProt Entry Name	APOM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/apolipoprotein-m-antibody-elisa/329145>