

Human Transferrin (TF) ELISA Kit | TF elisa kit

Human Transferrin (TF) ELISA Kit

CATALOG #
AAA330717

REACTIVITY
Human


PRODUCT SYNONYMS

Transferrin (TF); Human Transferrin (TF) ELISA Kit; TF; DKFZp781D0156; PRO1557; PRO2086; TF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	Human Transferrin (TF) ELISA Kit has high sensitivity and excellent specificity for detection of Human TF. No significant cross-reactivity or interference between Human TF and analogues was observed.
Sequence Length	698
Assay Type	Sandwich ELISA (Quantitative)
Samples	Cell Culture Supernatants, Serum, Plasma, Other Biological Fluids
Detection Method	Colorimetric
Usage Notes	* Do not mix components from different kit lots or use reagents beyond the kit expiration date. * Allow all reagents to warm to room temperature for at least 30 minutes before opening. * Pre-rinse the pipet tip with reagent, use fresh pipet tips for each sample, standard and reagent to avoid contamination. * Unused wells must be kept desiccated at 4 degree C in the sealed bag provided. * Mix Thoroughly is very important for the result. It is recommended using low frequency oscillator or slight hand shaking every 10 minutes. * It is recommended that all samples and standards be assayed in duplicate or triplicate.
Preparation and Storage	Store at 2-8 degree C.

IMAGES



Standard Curve (Sample)

Fig.1. Human Transferrin (TF) Standard Curve.

ADDITIONAL INFORMATION

Related Product Information for TF elisa kit	Description: This Human Transferrin (TF) ELISA Kit employs a two-site sandwich ELISA to quantitate TF in samples. An antibody specific for TF has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any TF present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TF is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TF bound in the initial step. The color development is stopped and the intensity of the color is measured. Background: Transferrin (TF) is a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of Transferrin (TF) is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. Transferrin (TF) may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	339453
NCBI GeneID	7018
UniProt Accession #	P02787
NCBI Official Full Name	transferrin
NCBI Official Synonym Full Names	transferrin
NCBI Official Symbol	TF
NCBI Official Synonym Symbols	TFQTL1; PRO1557; PRO2086; HEL-S-71p
NCBI Protein Information	serotransferrin
UniProt Protein Name	Serotransferrin
UniProt Gene Name	TF
UniProt Synonym Gene Names	Transferrin
UniProt Entry Name	TRFE_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/human-elisa-kits/transferrin-tf/330717>