

Mouse Tropomodulin-2 (TMOD2) ELISA Kit | TMOD2 elisa kit

Mouse Tropomodulin-2 (TMOD2) ELISA Kit

CATALOG #
AAA332167

REACTIVITY
Mouse


PRODUCT SYNONYMS

Tropomodulin-2 (TMOD2); Mouse Tropomodulin-2 (TMOD2) ELISA Kit; TMOD2; MGC39481; N-TMOD; NTMOD; neuronal tropomodulin; TMOD2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	Mouse Tropomodulin-2 (TMOD2) ELISA Kit has high sensitivity and excellent specificity for detection of Mouse TMOD2. No significant cross-reactivity or interference between Mouse TMOD2 and analogues was observed.
Sequence Length	351
Assay Type	Sandwich ELISA (Quantitative)
Samples	Cell Culture Supernatants, Serum, Plasma, Other Biological Fluids
Detection Method	Colorimetric
Usage Notes	* Do not mix components from different kit lots or use reagents beyond the kit expiration date. * Allow all reagents to warm to room temperature for at least 30 minutes before opening. * Pre-rinse the pipet tip with reagent, use fresh pipet tips for each sample, standard and reagent to avoid contamination. * Unused wells must be kept desiccated at 4 degree C in the sealed bag provided. * Mix Thoroughly is very important for the result. It is recommended using low frequency oscillator or slight hand shaking every 10 minutes. * It is recommended that all samples and standards be assayed in duplicate or triplicate.
Preparation and Storage	Store at 2-8 degree C.

IMAGES



Standard Curve (Sample)

Fig.1. Mouse Tropomodulin-2 (TMOD2) Standard Curve.

ADDITIONAL INFORMATION

Related Product Information for TMOD2 elisa kit	Description: This Mouse Tropomodulin-2 (TMOD2) ELISA Kit employs a two-site sandwich ELISA to quantitate TMOD2 in samples. An antibody specific for TMOD2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any TMOD2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TMOD2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TMOD2 bound in the initial step. The color development is stopped and the intensity of the color is measured. Background: TMOD2 encodes a neuronal-specific member of the tropomodulin family of actin-regulatory proteins. The encoded protein caps the pointed end of actin filaments preventing both elongation and depolymerization. The capping activity of this protein is dependent on its association with tropomyosin. Alternatively spliced transcript variants encoding different isoforms have been described. TMOD2 encodes a deduced 351-amino acid protein. Northern blot analysis demonstrated restricted expression of TMOD2 in neuronal tissues; an approximately 9.5-kb transcript was seen in all brain regions. Northern blot analysis showed that expression of mouse Tmod2 occurred as early as embryonic day 7 and progressively increased during development.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	84697031
NCBI GeneID	50876
NCBI Accession #	NP_001033799.1
NCBI GenBank Nucleotide #	NP_001033799.1
UniProt Accession #	Q9JKK7
NCBI Official Full Name	tropomodulin-2
NCBI Official Synonym Full Names	tropomodulin 2
NCBI Official Symbol	Tmod2
NCBI Official Synonym Symbols	NTMOD; N-Tmod
NCBI Protein Information	tropomodulin-2
UniProt Protein Name	Tropomodulin-2
UniProt Gene Name	Tmod2
UniProt Synonym Gene Names	N-Tmod
UniProt Entry Name	TMOD2_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.