

Mouse Tyrosine Hydroxylase (TH) ELISA Kit | TH elisa kit

Mouse Tyrosine Hydroxylase (TH) ELISA Kit

CATALOG #
AAA332385

REACTIVITY
Mouse


PRODUCT SYNONYMS

Tyrosine Hydroxylase (TH); Mouse Tyrosine Hydroxylase (TH) ELISA Kit; TH; DYT14; DYT5b; TYH; dystonia 14; tyrosine 3-monoxygenase; TH elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	Mouse Tyrosine hydroxylase (TH) ELISA Kit has high sensitivity and excellent specificity for detection of Mouse TH. No significant cross-reactivity or interference between Mouse TH and analogues was observed.
Sequence Length	498
Assay Type	Sandwich ELISA (Quantitative)
Samples	Cell Culture Supernatants, Serum, Plasma, Other Biological Fluids
Detection Method	Colorimetric
Usage Notes	* Do not mix components from different kit lots or use reagents beyond the kit expiration date. * Allow all reagents to warm to room temperature for at least 30 minutes before opening. * Pre-rinse the pipet tip with reagent, use fresh pipet tips for each sample, standard and reagent to avoid contamination. * Unused wells must be kept desiccated at 4 degree C in the sealed bag provided. * Mix Thoroughly is very important for the result. It is recommended using low frequency oscillator or slight hand shaking every 10 minutes. * It is recommended that all samples and standards be assayed in duplicate or triplicate.
Preparation and Storage	Store at 2-8 degree C.

IMAGES



Standard Curve (Sample)

Fig.1. Mouse Tyrosine hydroxylase (TH) Standard Curve.

ADDITIONAL INFORMATION

Related Product Information for TH elisa kit	Description: This Mouse Tyrosine hydroxylase (TH) ELISA Kit employs a two-site sandwich ELISA to quantitate TH in samples. An antibody specific for TH has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any TH present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TH is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TH bound in the initial step. The color development is stopped and the intensity of the color is measured. Background: The protein encoded by TH is involved in the conversion of tyrosine to dopamine. It is the rate-limiting enzyme in the synthesis of catecholamines, hence plays a key role in the physiology of adrenergic neurons. Mutations in TH have been associated with autosomal recessive Segawa syndrome. Alternatively spliced transcript variants encoding different isoforms have been noted for TH.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	6678337
NCBI GeneID	21823
NCBI Accession #	NP_033403.1
NCBI GenBank Nucleotide #	NP_033403.1
UniProt Accession #	P24529
NCBI Official Full Name	tyrosine 3-monooxygenase
NCBI Official Synonym Full Names	tyrosine hydroxylase
NCBI Official Symbol	Th
NCBI Protein Information	tyrosine 3-monooxygenase
UniProt Protein Name	Tyrosine 3-monooxygenase
UniProt Gene Name	Th
UniProt Synonym Gene Names	TH

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-elisa-kits/tyrosine-hydroxylase-th/332385>