

Human CCL2 ELISA Kit | CCL2 elisa kit

Human CCL2 ELISA Kit

CATALOG # AAA332530	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

CCL2; Human CCL2 ELISA Kit; GDCF-2; HC11; HSMCR30; MCAF; MCP-1; MCP1; MGC9434; SCYA2; SMC-CF; monocyte chemoattractant protein-1; monocyte chemotactic and activating factor; monocyte secretory protein JE; small inducible cytokine; CCL2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	Human CCL2 ELISA Kit has high sensitivity and excellent specificity for detection of HumanCCL2. No significant cross-reactivity or interference between Human CCL2 and analogues was observed.
Assay Type	Sandwich
Samples	Serum, Plasma, Cell culture supernatants
Detection Range	62.5 pg/mL-4,000 pg/mL
Sensitivity	30 pg/mL
Applicable Applications for CCL2 elisa kit	ELISA
Intra-assay Precision	Intra-assay Precision: The CV (%)?10%.
Inter-assay Precision	Inter-assay Precision :The CV (%)?12%
Preparation and Storage	The unopened kit should be stored at 2 - 8 degree C. After opening, please store refer to protocols.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CCL2 elisa kit	Principle of the Assay: CCL2 gene is one of several cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC subfamily of cytokines. Human CCL2 ELISA Kit employs a double antibody sandwich method to quantitate Human CCL2 in samples. An antibody specific for Human CCL2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any Human CCL2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for Human CCL2 is added to the wells. After washing, proprietary Streptavidin-HRP conjugates are added to the wells. Following a wash to remove any unbound streptavidin-enzyme reagent, adding HRP Substrate (TMB), TMB turns blue under the catalysis of HRP, and turns yellow after adding stop solution. Measure the OD value with a microplate reader at 450 nm wavelength. The Human CCL2 concentration is proportional to the OD450 nm value.
Product Categories/Family for CCL2 elisa kit	ELISA Assay Kit; Protein

NCBI AND UNIPROT INFORMATION

NCBI GI #	545465
NCBI GeneID	6347
UniProt Accession #	P13500; Q9UDF3; B2R4V3
Molecular Weight	11,025 Da
NCBI Official Full Name	MCP-1
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 2
NCBI Official Symbol	CCL2
NCBI Official Synonym Symbols	HC11; MCAF; MCP1; MCP-1; SCYA2; GDCF-2; SMC-CF; HSMCR30
NCBI Protein Information	C-C motif chemokine 2; small-inducible cytokine A2; monocyte secretory protein JE; monocyte chemotactic protein 1; monocyte chemoattractant protein 1; monocyte chemoattractant protein-1; monocyte chemotactic and activating factor; small inducible cytokine
UniProt Protein Name	C-C motif chemokine 2
UniProt Gene Name	CCL2
UniProt Synonym Gene Names	MCP1; SCYA2; MCAF; MCP-1
UniProt Entry Name	CCL2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/human-elisa-kits/ccl2/332530>