

Mouse IL-17 ELISA Kit | IL-17 elisa kit

Mouse IL-17 ELISA Kit

CATALOG # AAA332575	REACTIVITY Mouse	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

IL-17; Mouse IL-17 ELISA Kit; IL 17; IL17A; CTLA8; IL-17A; IL17; cytotoxic T-lymphocyte-associated antigen 8; cytotoxic T-lymphocyte-associated protein 8; cytotoxic T-lymphocyte-associated serine esterase 8; interleukin 17 (cytotoxic T-ly; IL-17 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	Mouse IL-17 ELISA Kit has high sensitivity and excellent specificity for detection of MouseIL-17. No significant cross-reactivity or interference between Mouse IL-17 and analogues was observed.
Assay Type	Sandwich
Samples	Serum, Plasma, Cell culture supernatants
Detection Range	7.82 pg/mL - 500 pg/mL
Sensitivity	4 pg/mL
Applicable Applications for IL-17 elisa kit	ELISA
Intra-assay Precision	Intra-assay Precision: The CV (%)?10%.
Inter-assay Precision	Inter-assay Precision :The CV (%)?12%
Preparation and Storage	The unopened kit should be stored at 2 - 8 degree C. After opening, please store refer to protocols.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL-17 elisa kit	Principle of the Assay: Interleukin 17 (IL-17; also known as IL-17A) is a 30-35 kDa variably glycosylated homodimeric protein that belongs to a uniquefamily of cysteine-knot related proteins. Its sequence was originally isolated from an activated rodent hybridoma and termedCTLA-8.Mouse IL-17 ELISA Kit employs a double antibody sandwich method to quantitate Mouse IL-17 in samples. Anantibody specific for Mouse IL-17 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells andany Mouse IL-17 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugatedantibody specific for Mouse IL-17 is added to the wells. After washing, proprietary Streptavidin-HRP conjugates is added to thewells. Following a wash to remove any unbound streptavidin-enzyme reagent, adding HRP Substrate (TMB), TMB turns blueunder the catalysis of HRP, and turns yellow after adding stop solution. Measure the OD value with a microplate reader at 450nmwavelength. The IL-17 concentration is proportional to the OD450 nm value.
Product Categories/Family for IL-17 elisa kit	ELISA Assay Kit; Protein

NCBI AND UNIPROT INFORMATION

NCBI GI #	6754324
NCBI GeneID	16171
NCBI Accession #	NP_034682.1
NCBI GenBank Nucleotide #	NM_010552.3
UniProt Accession #	Q62386; Q60971
Molecular Weight	17,490 Da
NCBI Official Full Name	interleukin-17A
NCBI Official Synonym Full Names	interleukin 17A
NCBI Official Symbol	Il17a
NCBI Official Synonym Symbols	Il17; Ctl8; IL-17; Ctl8; IL-17A
NCBI Protein Information	interleukin-17A; cytotoxic T-lymphocyte-associated antigen 8
UniProt Protein Name	Interleukin-17A
UniProt Gene Name	Il17a
UniProt Synonym Gene Names	Ctl8; Il17; IL-17; IL-17A; CTLA-8
UniProt Entry Name	IL17_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.