

Rabbit anti-Human, Mouse FASN Polyclonal Antibody | anti-FASN antibody

FASN Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA36634	Rabbit	Human, Mouse	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

FASN; Polyclonal Antibody; FASN Polyclonal Antibody; FAS; MGC14367; MGC15706; OA-519; SDR27X1; anti-FASN antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	2511
Applicable Applications for anti-FASN antibody	IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human FASN
Calculated Molecular Weight	272kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of A549 cells using FASN antibody. Blue: DAPI for nuclear staining.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse liver using FASN Antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver cancer using FASN Antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using FASN antibody at 1:200 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-FASN antibody

Fatty acid synthase (FASN) catalyzes the synthesis of long-chain fatty acids from acetyl-CoA and malonyl-CoA. FASN is active as a homodimer with seven different catalytic activities and produces lipids in the liver for export to metabolically active tissues or storage in adipose tissue. In most other human tissues, FASN is minimally expressed since they rely on circulating fatty acids for new structural lipid synthesis (1). Recently, increased expression of FASN has emerged as a phenotype common to most human carcinomas. In breast cancer, immunohistochemical staining showed that the levels of FASN are directly related to the size of breast tumors (2). Studies also showed that FASN is highly expressed in lung and prostate cancers and that FASN expression is an indicator of poor prognosis in breast and prostate cancer (3-5). Furthermore, inhibition of FASN is selectively cytotoxic to human cancer cells (5). Thus, increased interest has focused on FASN as a potential target for the identification and treatment of cancer as well as metabolic syndrome (6,7).

Product Categories/Family for anti-FASN antibody

Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	269849686
NCBI GeneID	2194
UniProt Accession #	P49327; Q13479; Q16702; Q4LE83; Q6P4U5; Q6SS02; Q969R1; Q96C68; Q96IT0
Molecular Weight	2511
NCBI Official Full Name	Fatty acid synthase
NCBI Official Synonym Full Names	fatty acid synthase
NCBI Official Symbol	FASN
NCBI Official Synonym Symbols	FAS; OA-519; SDR27X1
NCBI Protein Information	fatty acid synthase; short chain dehydrogenase/reductase family 27X, member 1
UniProt Protein Name	Fatty acid synthase
UniProt Gene Name	FASN
UniProt Synonym Gene Names	FAS
UniProt Entry Name	FAS_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/fasn-antibody-immunohistochemistry-western-blot-elisa/36634>