

Rabbit MAPK14-Y182-p Antibody | anti-MAPK14 antibody

Phospho-MAPK14-Y182-p Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA36726	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MAPK14-Y182-p; Phospho-MAPK14-Y182-p Polyclonal Antibody; Phospho-MAPK14-Y182: RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15 SAPK2A; p38ALPHA; anti-MAPK14 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Concentration	1mg/ml (varies by lot)
Sequence Length	360
Applicable Applications for anti-MAPK14 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic phosphorylated peptide around Y182 of human MAPK (NP_620581.1).
Preparation and Storage	Store at -20°C. Avoid freeze / thaw cycles.

IMAGES



WB (Western Blot)

Western blot analysis of 293T, using Phospho-p38 MAPK-Y182 antibody (AAA36726) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AAA36726) at 1:10000 dilution. Lysates/proteins: 25?g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK14 antibody	p38 MAP kinase (MAPK), also called RK (1) or CSBP (2), is the mammalian orthologue of the yeast HOG kinase that participates in a signaling cascade controlling cellular responses to cytokines and stress (1-4). Four isoforms of p38 MAPK, p38alpha, beta, gamma (also known as Erk6 or SAPK3), and delta (also known as SAPK4) have been identified. Similar to the SAPK/JNK pathway, p38 MAPK is activated by a variety of cellular stresses including osmotic shock, inflammatory cytokines, lipopolysaccharide (LPS), UV light, and growth factors (1-5). MKK3, MKK6, and SEK activate p38 MAPK by phosphorylation at Thr180 and Tyr182. Activated p38 MAPK has been shown to phosphorylate and activate MAPKAP kinase 2 (3) and to phosphorylate the transcription factors ATF-2 (5), Max (6), and MEF2 (5-8).SB203580 (4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)-imidazole) is a selective inhibitor of p38 MAPK. This compound inhibits the activation of MAPKAPK-2 by p38 MAPK and subsequent phosphorylation of HSP27 (9). SB203580 inhibits p38 MAPK catalytic activity by binding to the ATP-binding pocket, but does not inhibit phosphorylation of p38 MAPK by upstream kinases (10).
Product Categories/Family for anti-MAPK14 antibody	Phosphorylation

NCBI AND UNIPROT INFORMATION

NCBI GI #	2499600
NCBI GeneID	1432
UniProt Accession #	Q16539; O60776; Q13083; Q14084; Q8TDX0; A6ZJ92; A8K6P4; B0LPH0; B5TY32
Molecular Weight	Observed MW: 42kDa Calculated MW: 41kDa
NCBI Official Full Name	Mitogen-activated protein kinase 14
NCBI Official Synonym Full Names	mitogen-activated protein kinase 14
NCBI Official Symbol	MAPK14
NCBI Official Synonym Symbols	RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA
NCBI Protein Information	mitogen-activated protein kinase 14; MAP kinase 14; p38alpha Exp; p38 MAP kinase; MAP kinase Mxi2; MAP kinase p38 alpha; CSAID-binding protein; Csaids binding protein; MAX-interacting protein 2; stress-activated protein kinase 2A; p38 mitogen activated protein kinase; mitogen-activated protein kinase p38 alpha; cytokine suppressive anti-inflammatory drug binding protein; cytokine suppressive anti-inflammatory drug-binding protein
UniProt Protein Name	Mitogen-activated protein kinase 14
UniProt Gene Name	MAPK14
UniProt Synonym Gene Names	CSBP; CSBP1; CSBP2; CSPB1; MXI2; SAPK2A; MAP kinase 14; MAPK 14; CSAID-binding protein; CSBP; MAP kinase p38 alpha; SAPK2a
UniProt Entry Name	MK14_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.