

Rabbit IGF2BP3 Polyclonal Antibody | anti-IGF2BP3 antibody

IGF2BP3 Polyclonal Antibody

CATALOG # AAA36933	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Western Blot
------------------------------	-----------------------	--	-------------------------------------


PRODUCT SYNONYMS

IGF2BP3; Polyclonal Antibody; IGF2BP3 Polyclonal Antibody; KOC; CT98; IMP3; KOC1; IMP-3; VICKZ3; anti-IGF2BP3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	579
Applicable Applications for anti-IGF2BP3 antibody	WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human IGF2BP3
Calculated Molecular Weight	64kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human gastric cancer using IGF2BP3 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver using IGF2BP3 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat heart using IGF2BP3 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using IGF2BP3 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-IGF2BP3 antibody	The protein encoded by this gene is primarily found in the nucleolus, where it can bind to the 5' UTR of the insulin-like growth factor II leader 3 mRNA and may repress translation of insulin-like growth factor II during late development. The encoded protein contains several KH domains, which are important in RNA binding and are known to be involved in RNA synthesis and metabolism. A pseudogene exists on chromosome 7, and there are putative pseudogenes on other chromosomes.
Product Categories/Family for anti-IGF2BP3 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	254763311
NCBI GeneID	10643
UniProt Accession #	O00425; Q63HM0; Q6MZZ2; Q86VB1; A0A4Z5
Molecular Weight	579
NCBI Official Full Name	Insulin-like growth factor 2 mRNA-binding protein 3
NCBI Official Synonym Full Names	insulin-like growth factor 2 mRNA binding protein 3
NCBI Official Symbol	IGF2BP3
NCBI Official Synonym Symbols	KOC; CT98; IMP3; KOC1; IMP-3; VICKZ3
NCBI Protein Information	insulin-like growth factor 2 mRNA-binding protein 3; VICKZ family member 3; cancer/testis antigen 98; IGF2 mRNA-binding protein 3; IGF-II mRNA-binding protein 3; KH domain containing protein overexpressed in cancer
UniProt Protein Name	Insulin-like growth factor 2 mRNA-binding protein 3
UniProt Gene Name	IGF2BP3
UniProt Synonym Gene Names	IMP3; KOC1; VICKZ3; IGF2 mRNA-binding protein 3; IMP-3; hKOC
UniProt Entry Name	IF2B3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/igf2bp3-antibody-western-blot-elisa/36933>