

Rabbit anti-Human, Mouse C1QBP Polyclonal Antibody | anti-C1QBP antibody

C1QBP Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA36973	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

C1QBP; Polyclonal Antibody; C1QBP Polyclonal Antibody; p32; HABP1; gC1qR; GC1QBP; SF2p32; gC1Q-R; anti-C1QBP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	282
Applicable Applications for anti-C1QBP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human C1QBP
Calculated Molecular Weight	31kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cells using C1QBP antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using C1QBP antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-C1QBP antibody	C1QBP, also referred to as p32, p33, gC1q receptor (gC1qR), and hyaluronic acid binding protein 1 (HABP1), was originally identified via its binding interactions with Splicing Factor (SF-2) (1). Multiple, diverse binding partners of C1QBP were subsequently identified, including the globular heads of complement component C1q, hyaluronic acid, selected protein kinases (2), the tumor suppressor ARF (3-5), and multiple antigens of bacterial and viral origin (6). Research studies have shown that C1QBP is overexpressed in a number of cancer cell types (7), and has been implicated in the Warburg effect, whereby cancer cells shift their metabolism from oxidative phosphorylation to glycolysis (7). C1QBP has also been shown to inhibit the Mitochondrial Permeability Transition (MPT) pore, possibly serving a protective function against damage from oxidative stress (8).
Product Categories/Family for anti-C1QBP antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	730772
NCBI GeneID	708
UniProt Accession #	Q07021; Q2HXR8; Q9NNY8
Molecular Weight	282
NCBI Official Full Name	Complement component 1 Q subcomponent-binding protein, mitochondrial
NCBI Official Synonym Full Names	complement component 1, q subcomponent binding protein
NCBI Official Symbol	C1QBP
NCBI Official Synonym Symbols	p32; HABP1; gC1qR; GC1QBP; SF2p32; gC1Q-R
NCBI Protein Information	complement component 1 Q subcomponent-binding protein, mitochondrial; p33; glycoprotein gC1qBP; hyaluronan-binding protein 1; ASF/SF2-associated protein p32; mitochondrial matrix protein p32; C1q globular domain-binding protein; splicing factor SF2-associated protein
UniProt Protein Name	Complement component 1 Q subcomponent-binding protein, mitochondrial
UniProt Gene Name	C1QBP
UniProt Synonym Gene Names	GC1QBP; HABP1; SF2P32; C1qBP
UniProt Entry Name	C1QBP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/c1qbp-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/36973>