

Rabbit CFB Polyclonal Antibody | anti-CFB antibody

CFB Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA37036	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CFB; Polyclonal Antibody; CFB Polyclonal Antibody; BF; FB; BFD; GBG; CFAB; PBF2; AHUS4; FBI12; H2-Bf; anti-CFB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	764
Applicable Applications for anti-CFB antibody	IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human CFB
Calculated Molecular Weight	86kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse kidney using CFB antibody at dilution of 1:200 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CFB antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST.

ADDITIONAL INFORMATION

Related Product Information for anti-CFB antibody	This gene encodes complement factor B, a component of the alternative pathway of complement activation. Factor B circulates in the blood as a single chain polypeptide. Upon activation of the alternative pathway, it is cleaved by complement factor D yielding the noncatalytic chain Ba and the catalytic subunit Bb. The active subunit Bb is a serine protease which associates with C3b to form the alternative pathway C3 convertase. Bb is involved in the proliferation of preactivated B lymphocytes, while Ba inhibits their proliferation. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. This cluster includes several genes involved in regulation of the immune reaction. Polymorphisms in this gene are associated with a reduced risk of age-related macular degeneration. The polyadenylation site of this gene is 421 bp from the 5' end of the gene for complement component 2.
Product Categories/Family for anti-CFB antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	584908
NCBI GeneID	629
UniProt Accession #	P00751; O15006; Q29944; Q53F89; Q5JP67; Q5ST50; Q96HX6; Q9BTF5; Q9BX92; B0QZQ6
Molecular Weight	764
NCBI Official Full Name	Complement factor B
NCBI Official Synonym Full Names	complement factor B
NCBI Official Symbol	CFB
NCBI Official Synonym Symbols	BF; FB; BFD; GBG; CFAB; CFBD; PBF2; AHUS4; FBI12; H2-Bf; ARMD14
NCBI Protein Information	complement factor B; C3 proactivator; C3/C5 convertase; C3 proaccelerator; properdin factor B; B-factor, properdin; glycine-rich beta glycoprotein; glycine-rich beta-glycoprotein
UniProt Protein Name	Complement factor B
UniProt Gene Name	CFB
UniProt Synonym Gene Names	BF; BFD; GBG
UniProt Entry Name	CFAB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cfb-antibody-immunohistochemistry-western-blot-elisa/37036>