

Rabbit IL7 Polyclonal Antibody | anti-IL7 antibody

IL7 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA37194	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

IL7; Polyclonal Antibody; IL7 Polyclonal Antibody; IL-7; anti-IL7 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	177
Applicable Applications for anti-IL7 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human IL7
Calculated Molecular Weight	20kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IF (Immunofluorescence)
Immunofluorescence analysis of MCF-7 cells using IL7 antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using IL7 antibody at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-IL7 antibody	IL-7 plays a key role in lymphopoiesis and lymphoid homeostasis (1). Stromal and epithelial cells within the bone marrow and thymus produce IL-7 (1). The primary targets of IL-7 are T cells, B cells, and dendritic cells (1). IL-7 is crucial for T cell development, the importance of which is underscored by the lack of T cells in both mice and humans that are deficient in IL-7/IL-7R signaling (1,2). While IL-7 appears to be required for B cell development in mice, the role of IL-7 in human B cell development is unclear (1,3). In addition to its effects on T cell lymphopoiesis, IL-7 promotes the maintenance and survival of na ^{ve} and memory alphabeta T cells, as well as gammadelta T cells (1). The IL-7 receptor is a heterodimer of the common gamma chain, gammac, and the IL-7-specific IL-7Ralpha (1). IL-7 activates PI3K/Akt, Jak1/2, and Stat1, 3, and 5 (1).
Product Categories/Family for anti-IL7 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	124354
NCBI GeneID	3574
UniProt Accession #	P13232; Q5FBY5; Q5FBY9; A0N0L3
Molecular Weight	177
NCBI Official Full Name	Interleukin-7
NCBI Official Synonym Full Names	interleukin 7
NCBI Official Symbol	IL7
NCBI Official Synonym Symbols	IL-7
NCBI Protein Information	interleukin-7
UniProt Protein Name	Interleukin-7
UniProt Gene Name	IL7
UniProt Synonym Gene Names	IL-7
UniProt Entry Name	IL7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/il7-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/37194>