

Rabbit anti-Human CSTB Polyclonal Antibody | anti-CSTB antibody

CSTB Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA37447	Rabbit	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CSTB; Polyclonal Antibody; CSTB Polyclonal Antibody; PME; ULD; CST6; EPM1; STFB; EPM1A; anti-CSTB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	98
Applicable Applications for anti-CSTB antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human CSTB
Calculated Molecular Weight	11kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human prostate using CSTB antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CSTB antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-CSTB antibody	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and kininogens. This gene encodes a stefin that functions as an intracellular thiol protease inhibitor. The protein is able to form a dimer stabilized by noncovalent forces, inhibiting papain and cathepsins I, h and b. The protein is thought to play a role in protecting against the proteases leaking from lysosomes. Evidence indicates that mutations in this gene are responsible for the primary defects in patients with progressive myoclonic epilepsy (EPM1).
Product Categories/Family for anti-CSTB antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	1706278
NCBI GeneID	1476
UniProt Accession #	P04080
Molecular Weight	98
NCBI Official Full Name	Cystatin-B
NCBI Official Synonym Full Names	cystatin B (stefin B)
NCBI Official Symbol	CSTB
NCBI Official Synonym Symbols	PME; ULD; CST6; EPM1; STFB; EPM1A
NCBI Protein Information	cystatin-B; CPI-B; liver thiol proteinase inhibitor
UniProt Protein Name	Cystatin-B
UniProt Gene Name	CSTB
UniProt Synonym Gene Names	CST6; STFB
UniProt Entry Name	CYTB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cstb-antibody-flow-cytometry-immunofluorescence-immunohistochemistry-western-blot-elisa/37447>