

Rabbit H2AFZ Polyclonal Antibody | anti-H2AFZ antibody

H2AFZ Polyclonal Antibody

CATALOG # AAA37588	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

H2AFZ; Polyclonal Antibody; H2AFZ Polyclonal Antibody; H2AZ; H2A.z; H2A/z; H2A.Z-1; anti-H2AFZ antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	128
Applicable Applications for anti-H2AFZ antibody	IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human H2AFZ
Calculated Molecular Weight	13kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse liver using H2AFZ antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using H2AFZ antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse kidney using H2AFZ antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using H2AFZ antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-H2AFZ antibody

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent member of the histone H2A family that is distinct from other members of the family. Studies in mice have shown that this particular histone is required for embryonic development and indicate that lack of functional histone H2A leads to embryonic lethality.

Product Categories/Family for anti-H2AFZ antibody

Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	83288408
NCBI GeneID	3015
UniProt Accession #	P0C0S5; P17317; Q6I9U0; B2RD56
Molecular Weight	128
NCBI Official Full Name	Histone H2A.Z
NCBI Official Synonym Full Names	H2A histone family, member Z
NCBI Official Symbol	H2AFZ
NCBI Official Synonym Symbols	H2AZ; H2A.z; H2A/z; H2A.Z-1
NCBI Protein Information	histone H2A.Z; H2AZ histone
UniProt Protein Name	Histone H2A.Z
UniProt Gene Name	H2AFZ
UniProt Synonym Gene Names	H2AZ; H2A/z
UniProt Entry Name	H2AZ_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.