

Myostatin Active Protein | MSTN active protein

Recombinant Human Myostatin

CATALOG # AAA38198	HOST E Coli
------------------------------	-----------------------

PRODUCT SYNONYMS

Myostatin; Recombinant Human Myostatin; Myostatin Human; Myostatin Human Recombinant; GDF-8; MSTN; Growth Differentiation Factor 8; MSTN Muscle Hypertrophy; MSTN active protein

PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
Form/Format	Lyophilized from a concentrated (1mg/ml) solution containing no additives. Sterile Filtered White lyophilized (freeze-dried) powder.
Sequence	Thesequenc eofthefirs tfiveN-ter minalamino acidswasde terminedan dwasfoundt obeAsp-Phe -Gly-Leu-A sp
Sequence Length	375
Solubility	It is recommended to reconstitute the lyophilized Myostatin in sterile 20mM HCl at 0.1 mg/ml, which can then be further diluted to other aqueous solutions.
Protein Content	Protein quantitation was carried out by two independent methods:1. UV spectroscopy at 280 nm using the absorbency value of 1.55 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics). 2. Analysis by RP-HPLC, using a standard solution of Myostatin as a Reference Standard.
Biological Activity	The ED50 as determined by the inhibition of the proliferation of MPC-11 cells is < 20ng/ml, corresponding to a Specific Activity of 50,000units/mg.
Preparation and Storage	Lyophilized Myostatin although stable at room temperature for 3 weeks, should be stored desiccated below -18 degree C. Upon reconstitution Myostatin should be stored at 4 degree C between 2-7 days and for future use below -18 degree C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

ADDITIONAL INFORMATION

Related Product Information for MSTN active protein	Description: Myostatin Human Recombinant produced in E Coli is a homodimer, non-glycosylated polypeptide chain containing 2 x 109 amino acids and having a total molecular mass of 24814 Dalton. The GDF-8 is purified by proprietary chromatographic techniques. Introduction: GDF8 is a member of the bone morphogenetic protein (BMP) family and the TGF-beta superfamily. This group of proteins is characterized by a polybasic proteolytic processing site which is cleaved to produce a mature protein containing seven conserved cysteine residues. The members of this family are regulators of cell growth and differentiation in both embryonic and adult tissues. This gene is thought to encode a secreted protein which negatively regulates skeletal muscle growth.
Product Categories/Family for MSTN active protein	Cytokines and Growth Factors; Growth Factors; Myostatin

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885259
-----------	---------

NCBI GeneID	2660
NCBI Accession #	NP_005250.1
NCBI GenBank Nucleotide #	NM_005259.2
UniProt Accession #	O14793 ; Q6B0H2 ; A1C2J7 ; A1C2K0
Molecular Weight	42,750 Da
NCBI Official Full Name	growth/differentiation factor 8
NCBI Official Synonym Full Names	myostatin
NCBI Official Symbol	MSTN
NCBI Official Synonym Symbols	GDF8; MSLHP
NCBI Protein Information	growth/differentiation factor 8; GDF-8; growth differentiation factor 8
UniProt Protein Name	Growth/differentiation factor 8
UniProt Gene Name	MSTN
UniProt Synonym Gene Names	GDF8; GDF-8
UniProt Entry Name	GDF8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/e-coli-active-protein/myostatin/38198>