

Rabbit anti-Human DYRK2 Polyclonal Antibody | anti-DYRK2 antibody

DYRK2 Antibody

CATALOG # AAA39718	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunofluorescence, Immunocytochemistry, Western Blot, ELISA
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

DYRK2; Polyclonal Antibody; DYRK2 Antibody; Dual specificity tyrosine-phosphorylation-regulated kinase 2; dual-specificity tyrosine-(Y)-phosphorylation regulated kinase 2; anti-DYRK2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Two isoforms of DYRK2 are known to exist; this antibody will recognize both isoforms. DYRK2 antibody will not cross-react with other DYRK family members.
Purity/Purification	DYRK2 Antibody is affinity chromatography purified via peptide column.
Form/Format	Liquid
Concentration	1 mg/mL (varies by lot)
Sequence Length	528
Applicable Applications for anti-DYRK2 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot), ELISA
Conjugate	Unconjugated
Immunogen	DYRK2 antibody was raised against an 18 amino acid synthetic peptide near the carboxy terminus of human DYRK2.
Buffer	DYRK2 Antibody is supplied in PBS containing 0.02% sodium azide.
Preparation and Storage	DYRK2 antibody can be stored at 4 degree C for three months and -20 degree C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

IMAGES



IF (Immunofluorescence)
Immunofluorescence of DYRK2 in 293 cells with DYRK2 antibody at 20 µg/mL.

ICC (Immunocytochemistry)
Immunocytochemistry of DYRK2 in 293 cells with DYRK2 antibody at 10 µg/mL.



WB (Western Blot)
Western blot analysis of DYRK2 in 293 cell lysate with DYRK2 antibody at (A) 1 and (B) 2 µg/mL.

ADDITIONAL INFORMATION

Related Product Information for anti-DYRK2 antibody

DYRK2 Antibody: DYRK2 is a member of the dual-specificity tyrosine phosphorylation-regulated kinase (DYRK) family that is thought to be involved in cellular proliferation and apoptosis. Under normal conditions, nuclear but not cytoplasmic DYRK2 is ubiquitinated by MDM2, leading to its constitutive degradation. However, upon exposure to genotoxic stress, ATM phosphorylates DYRK2, leading to its dissociation from MDM2 and its phosphorylation of p53, thereby inducing apoptosis. Recent evidence also suggests that DYRK2 may serve as a scaffold that facilitates assembly of an E3 ubiquitin ligase.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503427
NCBI GeneID	8445
NCBI Accession #	NP_006473
NCBI GenBank Nucleotide #	NM_003583.3
UniProt Accession #	Q92630; Q9BRB5; B2R9V9
Molecular Weight	59,715 Da
NCBI Official Full Name	dual specificity tyrosine-phosphorylation-regulated kinase 2 isoform 1
NCBI Official Synonym Full Names	dual-specificity tyrosine-(Y)-phosphorylation regulated kinase 2
NCBI Official Symbol	DYRK2

NCBI Protein Information	dual specificity tyrosine-phosphorylation-regulated kinase 2
UniProt Protein Name	Dual specificity tyrosine-phosphorylation-regulated kinase 2
UniProt Gene Name	DYRK2
UniProt Entry Name	DYRK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/dyrk2-antibody-immunofluorescence-immunocytochemistry-western-blot-elisa/39718>