

## Rabbit CCL17 Polyclonal Antibody | anti-CCL17 antibody

CCL17 Antibody

| CATALOG # | HOST   | REACTIVITY        | APPLICATIONS                                                  |
|-----------|--------|-------------------|---------------------------------------------------------------|
| AAA40799  | Rabbit | Human, Mouse, Rat | Immunofluorescence, Immunohistochemistry, Western Blot, ELISA |


### PRODUCT SYNONYMS

CCL17; Polyclonal Antibody; CCL17 Antibody; TARC; ABCD-2; SCYA17; A-152E5.3; C-C motif chemokine 17; CC chemokine TARC; chemokine (C-C motif) ligand 17; anti-CCL17 antibody

### PRODUCT SPECIFICATIONS


|                                                 |                                                                                                                       |
|-------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------|
| Host                                            | Rabbit                                                                                                                |
| Reactivity                                      | Human, Mouse, Rat                                                                                                     |
| Clonality                                       | Polyclonal                                                                                                            |
| Isotype                                         | IgG                                                                                                                   |
| Purity/Purification                             | CCL17 Antibody is affinity chromatography purified via peptide column.                                                |
| Form/Format                                     | Liquid                                                                                                                |
| Concentration                                   | 1 mg/mL (varies by lot)                                                                                               |
| Sequence Length                                 | 94                                                                                                                    |
| Applicable Applications for anti-CCL17 antibody | IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA                                         |
| Conjugate                                       | Unconjugated                                                                                                          |
| Immunogen                                       | Rabbit polyclonal CCL17 antibody was raised against a 16 amino acid peptide near the carboxy terminus of human CCL17. |
| Buffer                                          | CCL17 Antibody is supplied in PBS containing 0.02% sodium azide.                                                      |
| Preparation and Storage                         | CCL17 antibody can be stored at 4 degree C for three months and -20 degree C, stable for up to one year.              |

### IMAGES




#### IF (Immunofluorescence)

Immunofluorescence of CCL17 in human spleen tissue with CCL17 antibody at 20 µg/mL.



#### IHC (Immunohistochemistry)

Immunohistochemistry of CCL17 in human spleen tissue with CCL17 antibody at 5 µg/mL.



#### WB (Western Blot)

Western blot analysis of CCL17 in rat spleen tissue lysate with CCL17 antibody at (A) 1 and (B) 2 &#956;g/mL

### ADDITIONAL INFORMATION

#### Related Product Information for anti-CCL17 antibody

CCL17 Antibody: CCL17, also known as TARC, binds to chemokine receptors CCR4 and CCR8 and plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells. CCL17 is secreted from monocyte-derived DCs and endothelial cells and is responsible for selective recruitment and migration of activated Th2 lymphocytes. CCL17 is also required for CCR7 and CXCR4-dependent migration of cutaneous dendritic cells under inflammatory conditions.

### NCBI AND UNIPROT INFORMATION

|                                  |                                                                                                                                                                                                             |
|----------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NCBI GI #                        | 4506829                                                                                                                                                                                                     |
| NCBI GeneID                      | 6361                                                                                                                                                                                                        |
| NCBI Accession #                 | NP_002978                                                                                                                                                                                                   |
| NCBI GenBank Nucleotide #        | NM_002987.2                                                                                                                                                                                                 |
| UniProt Accession #              | Q92583; Q2M287; A0N0Q9                                                                                                                                                                                      |
| Molecular Weight                 | 10 kDa                                                                                                                                                                                                      |
| NCBI Official Full Name          | C-C motif chemokine 17                                                                                                                                                                                      |
| NCBI Official Synonym Full Names | chemokine (C-C motif) ligand 17                                                                                                                                                                             |
| NCBI Official Symbol             | CCL17                                                                                                                                                                                                       |
| NCBI Official Synonym Symbols    | TARC; ABCD-2; SCYA17; A-152E5.3                                                                                                                                                                             |
| NCBI Protein Information         | C-C motif chemokine 17; CC chemokine TARC; T cell-directed CC chemokine; small-inducible cytokine A17; thymus and activation-regulated chemokine; small inducible cytokine subfamily A (Cys-Cys), member 17 |
| UniProt Protein Name             | C-C motif chemokine 17                                                                                                                                                                                      |
| UniProt Gene Name                | CCL17                                                                                                                                                                                                       |
| UniProt Synonym Gene Names       | SCYA17; TARC                                                                                                                                                                                                |
| UniProt Entry Name               | CCL17_HUMAN                                                                                                                                                                                                 |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)

Online datasheet: <https://www.aaabiotech.com/polyclonal/ccl17-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/40799>