

Rat Cystatin C ELISA Kit | CST3 elisa kit

Rat Cystatin C PicoKine ELISA Kit

CATALOG #
AAA45856

REACTIVITY
Rat


PRODUCT SYNONYMS

Cystatin C; Rat Cystatin C PicoKine ELISA Kit; Cystatin-C; cystatin C; Cst3; Cystatin-3; CYTC_HUMAN; Gamma-trace; Neuroendocrine basic polypeptide; Post-gamma-globulin; CST3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	Natural and recombinant rat Cystatin C
Sequence Length	140
Samples	Cell culture supernates, serum, plasma (heparin, EDTA) and urine
Assay Type	Sandwich
Detection Range	312pg/ml-20,000pg/ml
Sensitivity	< 10pg/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay) Three samples of known concentration were tested on one plate to assess intra-assay precision.
Inter-assay Precision	Inter-Assay Precision (Precision between assays) Three samples of known concentration were tested in separate assays to assess inter-assay precision.
Preparation and Storage	Store at 4 degree C for 6 months, at -20 degree C for 12 months. Avoid multiple freeze-thaw cycles (Shipped with wet ice.)

IMAGES



Standard Curve (Sample)

Rat Cystatin C PicoKine ELISA Kit standard curve

ADDITIONAL INFORMATION

Related Product Information for CST3 elisa kit	Principle of the assay: rat Cystatin C ELISA Kit was based on standard sandwich enzyme-linked immune-sorbent assay technology. A monoclonal antibody from mouse specific for Cystatin C has been precoated onto 96-well plates. Standards(NSO, M1-A140) and test samples are added to the wells, a biotinylated detection polyclonal antibody from goat specific for Cystatin C is added subsequently and then followed by washing with PBS or TBS buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with PBS or TBS buffer. HRP substrate TMB was used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the rat Cystatin C amount of sample captured in plate. Background: Cystatin C or cystatin 3 (formerly gamma trace, post-gamma-globulin or neuroendocrine basic polypeptide), a protein encoded by the CST3 gene, was originally described as a constituent of normal cerebrospinal fluid (CSF) and of urine from patients with renal failure. Cystatin 3 has a low molecular weight (approximately 13.3 kilodaltons), and it is removed from the bloodstream by glomerular filtration in the kidneys. In mice, all cells with a nucleus (cell core containing the DNA) produce cystatin C as a chain of 120 amino acids. It is found in virtually all tissues and bodily fluids. Cystatin C, which belongs to the type II cystatin gene family, is a potent inhibitor of lysosomal proteinases (enzymes from a special subunit of the cell that break down proteins) and probably one of the most important extracellular inhibitors of cysteine proteases (it prevents the breakdown of proteins outside the cell by a specific type of protein degrading enzymes). Moreover, cystatin C is involved in network reorganization in the epileptic dentate gyrus.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	83301921
NCBI GeneID	25307
UniProt Accession #	P14841; Q5M968
Molecular Weight	15,437 Da
NCBI Official Full Name	Cystatin-C
NCBI Official Synonym Full Names	cystatin C
NCBI Official Symbol	Cst3
NCBI Official Synonym Symbols	CYSC
NCBI Protein Information	cystatin-C; cystatin 3; cystatin-3; Cystatin C (cysteine proteinase inhibitor)
UniProt Protein Name	Cystatin-C
UniProt Gene Name	Cst3
UniProt Entry Name	CYTC_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.