

Rabbit anti-Human PMEL17/GP100 Monoclonal Antibody | anti-PMEL antibody

Anti-PMEL17/GP100 Rabbit Monoclonal Antibody

CATALOG # AAA47044	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

PMEL17/GP100; Monoclonal Antibody; Anti-PMEL17/GP100 Rabbit Monoclonal Antibody; Melanocyte protein PMEL; ME20-M; ME20M; Melanocyte protein Pmel 17; Melanocytes lineage-specific antigen GP100; Melanoma-associated ME20 antigen; P1; P100; Premelanosome protein; Silver locus protein homolog; M-alpha; 95 kDa melanocyte-specific secreted glycoprotein; P26; Secreted melanoma-associated ME20 antigen; ME20-S; ME20S; M-beta; PMEL; D12S53E; PMEL17; SILV; anti-PMEL antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	626
Applicable Applications for anti-PMEL antibody	WB (Western Blot), IHC (Immunohistochemistry)
Immunogen	A synthesized peptide derived from human PMEL17/GP100
Subcellular Localization	Endoplasmic reticulum membrane; Single-pass type I membrane protein. Golgi apparatus. Melanosome. Endosome, multivesicular body. Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Localizes predominantly to intraluminal vesicles (ILVs) within multivesicular bodies. Associates with ILVs found within the lumen of premelanosomes and melanosomes and particularly in compartments that serve as precursors to the striated stage II premelanosomes.
Tissue Specificity	Preferentially expressed in melanomas. Some expression was found in dysplastic nevi. Not found in normal tissues nor in carcinomas. Normally expressed at low levels in quiescent adult melanocytes but overexpressed by proliferating neonatal melanocytes and during tumor growth.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human melanoma, using PMEL17 / GP100 Antibody(AAA47044)PMEL was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-PMEL Antibody (AAA47044)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)
Western blot analysis of PMEL17 / GP100 expression in human melanoma lysate (AAA47044).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PMEL monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PMEL

ADDITIONAL INFORMATION

Related Product Information for anti-PMEL antibody	Plays a central role in the biogenesis of melanosomes. Involved in the maturation of melanosomes from stage I to II. The transition from stage I melanosomes to stage II melanosomes involves an elongation of the vesicle, and the appearance within of distinct fibrillar structures. Release of the soluble form, ME20-S, could protect tumor cells from antibody mediated immunity.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	318037593
NCBI GeneID	6490
NCBI Accession #	NP_001186982.1
NCBI GenBank Nucleotide #	NM_001200053.1
UniProt Accession #	P40967; Q12763; Q14448; Q14817; Q16565; B3KS57; B7Z6D7
Molecular Weight	70255 MW
NCBI Official Full Name	melanocyte protein PMEL isoform 2
NCBI Official Synonym Full Names	premelanosome protein
NCBI Official Symbol	PMEL
NCBI Official Synonym Symbols	P1; SI; SIL; ME20; P100; SILV; ME20M; gp100; ME20-M; PMEL17; D12S53E
NCBI Protein Information	melanocyte protein PMEL
UniProt Protein Name	Melanocyte protein PMEL
UniProt Gene Name	PMEL
UniProt Synonym Gene Names	D12S53E; PMEL17; SILV; ME20M; ME20-S; ME20S

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/monoclonal/pm17-gp100-antibody-western-blot-immunohistochemistry-elisa/47044>