

CD22 recombinant protein

Recombinant human CD22 protein with C-terminal human Fc and 6xHis tag

CATALOG # AAA47189	HOST HEK293 cells	APPLICATIONS Western Blot
------------------------------	-----------------------------	-------------------------------------

PRODUCT SYNONYMS

CD22; Recombinant human CD22 protein with C-terminal human Fc and 6xHis tag; Human CD22 Protein; hFc-His Tag; SIGLEC2; BL-CAM; SIGLEC-2; Siglec2; SIGLEC2FLJ22814; CD22 recombinant protein

PRODUCT SPECIFICATIONS


Host	HEK293 cells
Purity/Purification	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Form/Format	Lyophilized from sterile PBS, pH 7.4. Normally 5%-12% trehalose is added as protectants before lyophilization.
Sequence	CD22(Asp20 -Arg687)+h Fc(Glu99-A la330)
Applicable Applications for CD22 recombinant protein	WB (Western Blot)
Source	Human
Conjugation	C-Human Fc and 6xHis tag
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -80 degree C for 12 months (Avoid repeated freezing and thawing) Shipping: Blue Ice

IMAGES



SDS-PAGE

Figure 1. Human CD22, hFc-His Tag on SDS-PAGE under reducing condition.



ELISA

Figure 1. Human CD22, hFc-His Tag on SDS-PAGE under reducing condition.

ADDITIONAL INFORMATION

Related Product Information for CD22 recombinant protein	B-cell receptor CD22 is also known as Sialic acid-binding Ig-like lectin 2 (Siglec-2), B-lymphocyte cell adhesion molecule (BL-CAM), T-cell surface antigen Leu-14, which belongs to the immunoglobulin superfamily and SIGLEC (sialic acid binding Ig-like lectin) family. CD22 mediates B-cell B-cell interactions, and may be involved in the localization of B-cells in lymphoid tissues. Siglec-2/CD22 binds sialylated glycoproteins, one of which is CD45. Siglec2/CD22 plays a role in positive regulation through interaction with Src family tyrosine kinases and may also act as an inhibitory receptor by recruiting cytoplasmic phosphatases via their SH2 domains that block signal transduction through dephosphorylation of signaling molecules.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	297374825
NCBI GeneID	933
NCBI Accession #	NP_001172028.1
NCBI GenBank Nucleotide #	NM_001185099.1
UniProt Accession #	P20273; O95699; O95701; O95702; O95703; Q01665; Q92872; Q92873; Q9UQA6; Q9UQA7; Q9UQA8; F5H7U3
Molecular Weight	95,348 Da
NCBI Official Full Name	B-cell receptor CD22 isoform 2
NCBI Official Synonym Full Names	CD22 molecule
NCBI Official Symbol	CD22
NCBI Official Synonym Symbols	SIGLEC2; SIGLEC-2
NCBI Protein Information	B-cell receptor CD22; BL-CAM; CD22 antigen; T-cell surface antigen Leu-14; B-lymphocyte cell adhesion molecule; sialic acid binding Ig-like lectin 2; sialic acid-binding Ig-like lectin 2
UniProt Protein Name	B-cell receptor CD22
UniProt Gene Name	CD22
UniProt Synonym Gene Names	SIGLEC2; BL-CAM; Siglec-2
UniProt Entry Name	CD22_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.