

CD40 recombinant protein

Recombinant human CD40 protein with C-terminal mouse Fc and 6xHis tag

CATALOG # AAA47197	HOST HEK293 cells	APPLICATIONS Western Blot
------------------------------	-----------------------------	-------------------------------------

PRODUCT SYNONYMS

CD40; Recombinant human CD40 protein with C-terminal mouse Fc and 6xHis tag; Human CD40 Protein; mFc-His Tag; Bp50; CDW40; MGC9013; TNFRSF5; p50; CD40 recombinant protein

PRODUCT SPECIFICATIONS

Host	HEK293 cells
Purity/Purification	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Form/Format	Lyophilized from sterile PBS, pH 7.4. Normally 5%-22% trehalose is added as protectants before lyophilization.
Sequence	CD40(Glu21 -Arg193)+m Fc(Pro99-L ys330)
Applicable Applications for CD40 recombinant protein	WB (Western Blot)
Source	Human
Conjugation	C-Mouse Fc and 6xHis tag
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -80 degree C for 12 months (Avoid repeated freezing and thawing) Shipping: Blue Ice

IMAGES


SDS-PAGE

Figure 1. Human CD40 Protein, mFc-His Tag on SDS-PAGE under reducing condition.



ELISA

Figure 1. Human CD40 Protein, mFc-His Tag on SDS-PAGE under reducing condition.

ADDITIONAL INFORMATION

Related Product Information for CD40 recombinant protein	This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Mutations affecting this gene are the cause of autosomal recessive hyper-IgM immunodeficiency type 3 (HIGM3). Multiple alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507581
NCBI GeneID	958
NCBI Accession #	NP_001241.1
NCBI GenBank Nucleotide #	NM_001250.4
UniProt Accession #	P25942; Q53GN5; Q5JY15; Q5U007; Q7M4Q8; Q86YK5; Q9BYU0; E1P5S9
Molecular Weight	30,619 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 5 isoform 1
NCBI Official Synonym Full Names	CD40 molecule, TNF receptor superfamily member 5
NCBI Official Symbol	CD40
NCBI Official Synonym Symbols	p50; Bp50; CDW40; TNFRSF5
NCBI Protein Information	tumor necrosis factor receptor superfamily member 5; CD40L receptor; CD40 type II isoform; B cell-associated molecule; B cell surface antigen CD40; B-cell surface antigen CD40; CD40 antigen (TNF receptor superfamily member 5); tumor necrosis factor recept
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 5
UniProt Gene Name	CD40
UniProt Synonym Gene Names	TNFRSF5
UniProt Entry Name	TNR5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/cd40-recombinant-protein-hek293-wb-western-blot/47197>