

NKG2D recombinant protein

Recombinant human NKG2D protein with N-terminal mouse Fc

CATALOG # AAA47239	HOST HEK293 cells	APPLICATIONS Western Blot
------------------------------	-----------------------------	-------------------------------------

PRODUCT SYNONYMS

NKG2D; Recombinant human NKG2D protein with N-terminal mouse Fc; Human NKG2D Protein; mFc Tag; NKG2D;CD314;KLRK1;NK cell receptor D; NKG2D recombinant protein

PRODUCT SPECIFICATIONS

Host	HEK293 cells
Purity/Purification	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Form/Format	Lyophilized from sterile PBS, pH 7.4. Normally 5%-68% trehalose is added as protectants before lyophilization.
Sequence	mFc(Pro99- Lys330)+NK G2D(Ile73- Val216)
Applicable Applications for NKG2D recombinant protein	WB (Western Blot)
Source	Human
Conjugation	N-Mouse Fc
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -80 degree C for 12 months (Avoid repeated freezing and thawing) Shipping: Blue Ice

IMAGES


SDS-PAGE

Figure 1. Human NKG2D Protein, mFc Tag on SDS-PAGE under reducing condition.



ELISA

Figure 1. Human NKG2D Protein, mFc Tag on SDS-PAGE under reducing condition.

ADDITIONAL INFORMATION

Related Product Information for NKG2D recombinant protein	Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of NK and T cells. The surface expression of these ligands is important for the recognition of stressed cells by the immune system, and thus this protein and its ligands are therapeutic targets for the treatment of immune diseases and cancers. Read-through transcription exists between this gene and the upstream KLRC4 (killer cell lectin-like receptor subfamily C, member 4) family member in the same cluster.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	315221164
NCBI GeneID	100528032
NCBI Accession #	NP_001186734.1
NCBI GenBank Nucleotide #	NM_001199805.1
UniProt Accession #	P26718; Q9NR41; A8K7K5; A8K7P4
Molecular Weight	25,274 Da
NCBI Official Full Name	NKG2-D type II integral membrane protein
NCBI Official Symbol	KLRC4-KLRK1
NCBI Protein Information	NKG2-D type II integral membrane protein; NK cell receptor D; NKG2-D-activating NK receptor; killer cell lectin-like receptor subfamily K member 1
UniProt Protein Name	NKG2-D type II integral membrane protein
UniProt Gene Name	KLRK1
UniProt Synonym Gene Names	D12S2489E; NKG2D
UniProt Entry Name	NKG2D_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/klrk1-recombinant-protein-hek293-wb-western-blot/47239>