

Rabbit anti-Human CD27 Monoclonal Antibody | anti-CD27 antibody

Anti-CD27 antibody (DM59), Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA47297	Rabbit	Human	Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

CD27; Monoclonal Antibody; Anti-CD27 antibody (DM59); Rabbit mAb; TNFRSF7; S152; T14; Tp55; anti-CD27 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	DM59
Purity/Purification	Purified from cell culture supernatant by affinity chromatography
Form/Format	Liquid. 50% Glycerol; PBS, pH7.4; 0.1% BSA; 0.1% Procline 300
Applicable Applications for anti-CD27 antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human CD27 (Ala20-Arg191) produced by using human HEK293 cells
Reconstitution	Reconstitute with deionized water
Preparation and Storage	Store at -20 degree C for 12 months (Avoid repeated freezing and thawing)

IMAGES



FCM/FACS (Flow Cytometry)

Figure 1. Expi 293 cell line transfected with irrelevant protein (A) and human CD27 (B) were surface stained with Rabbit anti-CD27 monoclonal antibody 1µg/ml (clone: DM59) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.



FCM/FACS (Flow Cytometry)

Figure 1. Expi 293 cell line transfected with irrelevant protein (A) and human CD27 (B) were surface stained with Rabbit anti-CD27 monoclonal antibody 1µg/ml (clone: DM59) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CD27 antibody

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507587
NCBI GeneID	939
NCBI Accession #	NP_001233.1
NCBI GenBank Nucleotide #	NM_001242.4
UniProt Accession #	P26842; B2RDZ0
Molecular Weight	29,137 Da
NCBI Official Full Name	CD27 antigen
NCBI Official Synonym Full Names	CD27 molecule
NCBI Official Symbol	CD27
NCBI Official Synonym Symbols	T14; S152; Tp55; TNFRSF7; S152. LPFS2
NCBI Protein Information	CD27 antigen; CD27L receptor; T cell activation antigen S152; T-cell activation antigen CD27; tumor necrosis factor receptor superfamily, member 7
UniProt Protein Name	CD27 antigen
UniProt Gene Name	CD27
UniProt Synonym Gene Names	TNFRSF7
UniProt Entry Name	CD27_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/cd27-antibody-clone-dm59-flow-cytometry-elisa/47297>