

Mouse CD221 Monoclonal Antibody | anti-CD221 antibody

MOUSE ANTI HUMAN CD221:RPE

CATALOG # AAA49498	HOST Mouse	APPLICATIONS Flow Cytometry, Functional Assay
------------------------------	----------------------	---


PRODUCT SYNONYMS

CD221; Monoclonal Antibody; MOUSE ANTI HUMAN CD221:RPE; anti-CD221 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	1H7
Form/Format	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilised
Sequence Length	1,367
Applicable Applications for anti-CD221 antibody	FCM/FACS (Flow Cytometry)
Reconstitution	Reconstitute with 1 ml distilled water Perservative Stabilisers
Immunogen	Purified human placental IGF-I receptor Fusion Partners
Target Species	Human
Preparation and Storage	Prior to reconstitution store at 4 degree C. Following reconstitution store at 4 degree C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 12 months from date of reconstitution.

IMAGES




Application Data

Staining of human peripheral blood granulocytes with Mouse anti Human CD221




Application Data

Staining of human peripheral blood granulocytes with Mouse anti Human CD221: Low Endotoxin



Application Data

Staining of human peripheral blood granulocytes with Mouse anti Human CD221: Alexa Fluor 488



Application Data

Staining of human peripheral blood granulocytes with Mouse anti Human CD221: RPE

ADDITIONAL INFORMATION

Related Product Information for anti-CD221 antibody

Mouse anti Human CD221 antibody, clone 1H7 recognizes human CD221, a 155kD receptor tyrosine kinase, also known as Insulin-like growth factor I receptor (IGF-I Receptor). CD221 is composed of two extracellular alpha-subunits and two transmembrane beta-subunits. Clone 1H7 recognizes an epitope in the alpha subunits of CD221, demonstrated by Western blotting (Li et al.1993). CD221 is expressed in a range of tissues, including kidney, liver, placenta, mammary gland, brain, ovary and skin. The ligands for CD221 include IGF-I and IGF-II, which bind to CD221 and activate tyrosine kinase activity, resulting in phosphorylation of several intracellular signalling proteins. Clone 1H7 is reported to partially block binding of IGF-I and IGF-II to CD221 (Li et al.1993).

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557665
NCBI GeneID	3480
NCBI Accession #	NP_000866.1
NCBI GenBank Nucleotide #	NM_000875.4
UniProt Accession #	P08069; Q14CV2; Q9UCC0; B1B5Y2
Molecular Weight	154,793 Da
NCBI Official Full Name	insulin-like growth factor 1 receptor isoform 1
NCBI Official Synonym Full Names	insulin-like growth factor 1 receptor
NCBI Official Symbol	IGF1R
NCBI Official Synonym Symbols	IGFR; CD221; IGFR; JTK13
NCBI Protein Information	insulin-like growth factor 1 receptor; IGF-I receptor; insulin-like growth factor I receptor; soluble IGF1R variant 1; soluble IGF1R variant 2
UniProt Protein Name	Insulin-like growth factor 1 receptor
UniProt Gene Name	IGF1R
UniProt Synonym Gene Names	IGF-I receptor
UniProt Entry Name	IGF1R_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.