

Rabbit anti-Mouse 4EBP1 Polyclonal Antibody | anti-4EBP1 antibody

RABBIT ANTI 4EBP1 (C-TERMINAL)


CATALOG # AAA49986	HOST Rabbit	REACTIVITY Mouse	APPLICATIONS Western Blot
------------------------------	-----------------------	----------------------------	-------------------------------------

PRODUCT SYNONYMS


4EBP1; Polyclonal Antibody; RABBIT ANTI 4EBP1 (C-TERMINAL); anti-4EBP1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Isotype	IgG
Specificity	This item recognises an epitope within the C-terminal region (CT) of eukaryotic translation initiation factor 4E binding protein 1 (4EBP1), a rapamycin-sensitive downstream target of TOR (Target of rapamycin), and regulatory factor of eukaryotic mRNA translation. The association between 4EBP1 and eukaryotic translation initiation factor 4E (eIF-4E) inhibits the binding of eIF-4E with p220, thereby preventing the formation of a functional eIF-4E complex, essential for mRNA translation. Phosphorylation of 4EBP1 following stimulation by mitogens such as insulin, results in the release of eIF-4E from 4EBP1, thereby enabling the translation complex to assemble. This item has been reported as suitable for use in IHC on mouse 3T3 cell cytopins.
Form/Format	Purified Purified IgG - liquid
Concentration	IgG concentration 1.0mg/ml (varies by lot)
Sequence Length	118
Applicable Applications for anti-4EBP1 antibody	WB (Western Blot)
Perservative Stabilisers	0.02% Sodium Azide (NaN ₃) Preparation
Immunogen	A 14 amino acid peptide located near human 4EBP1 carboxy-terminus. Buffer Solution
Target Species	Human
Preparation and Storage	Store at 4 degree C or at -20 degree C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 18 months from date of despatch.



Application Data
Immunocytochemical staining of 3T3 cells with Rabbit anti Human 4E-BP1 (AAA49986)



Application Data
Western blot analysis of whole cell lysates from 3T3 mouse embryonic fibroblast cells probed with Rabbit anti Human 4E-BP1 (AAA49986) at 2.5(A), 5(B) and 10(C) ug/ml

ADDITIONAL INFORMATION

Related Product Information for anti-4EBP1 antibody	MBS221870 recognises an epitope within the C-terminal region (CT) of eukaryotic translation initiation factor 4E binding protein 1 (4EBP1), a rapamycin-sensitive downstream target of TOR (Target of rapamycin), and regulatory factor of eukaryotic mRNA translation. The association between 4EBP1 and eukaryotic translation initiation factor 4E (eIF-4E) inhibits the binding of eIF-4E with p220, thereby preventing the formation of a functional eIF-4E complex, essential for mRNA translation. Phosphorylation of 4EBP1 following stimulation by mitogens such as insulin, results in the release of eIF-4E from 4EBP1, thereby enabling the translation complex to assemble. MBS221870 has been reported as suitable for use in IHC on mouse 3T3 cell cytopins.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758258
NCBI GeneID	1978
NCBI Accession #	NP_004086.1
NCBI GenBank Nucleotide #	NM_004095.3
UniProt Accession #	Q13541; Q6IBN3; B2R502; D3DSW8
Molecular Weight	12,580 Da
NCBI Official Full Name	eukaryotic translation initiation factor 4E-binding protein 1
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 4E binding protein 1
NCBI Official Symbol	EIF4EBP1
NCBI Official Synonym Symbols	BP-1; 4EBP1; 4E-BP1; PHAS-I
NCBI Protein Information	eukaryotic translation initiation factor 4E-binding protein 1; eIF4E-binding protein 1; phosphorylated heat- and acid-stable protein regulated by insulin 1
UniProt Protein Name	Eukaryotic translation initiation factor 4E-binding protein 1
UniProt Gene Name	EIF4EBP1
UniProt Synonym Gene Names	4E-BP1; eIF4E-binding protein 1; PHAS-I
UniProt Entry Name	4EBP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

