

Rabbit anti-Human c-Met Polyclonal Antibody | anti-MET antibody

Rabbit Polyclonal to Human c-Met

CATALOG # AAA51007	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

c-Met; Polyclonal Antibody; Rabbit Polyclonal to Human c-Met; Anti-c-Met Antibody (C-Terminus) IHC-plus; AUTS9; C-Met; HGF/SF receptor; HGFR; Met kinase; Proto-oncogene c-Met; Scatter factor receptor; SF receptor; HGF receptor; RCCP2; Tyrosine-protein kinase Met; Human c-Met; anti-MET antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Immunoaffinity Purified
Form/Format	10mM PBS/1% BSA buffer pH 7.4 with 0.1% sodium azide.
Concentration	0.2 mg/ml (varies by lot)
Sequence Length	1390
Applicable Applications for anti-MET antibody	IHC (Immunohistochemistry)
Target Species	Human
Immunogen Description	A synthetic peptide from C-terminus of human c-Met protein.
Immunogen Type	Synthetic peptide
Immunogen	c-Met antibody was raised against a synthetic peptide from C-terminus of human c-Met protein.
Antigen Modification	C-Terminus
Preparation and Storage	Store at 2 degree C to 8 degree C degrees. Do not freeze.

IMAGES



IHC (Immunohistochemistry)

Anti-c-Met antibody IHC of human liver. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 15 ug/ml.



IHC (Immunohistochemistry)

Anti-c-Met antibody IHC of human placenta. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 15 ug/ml.

ADDITIONAL INFORMATION

Related Product Information for anti-MET antibody	Met, an HGF Receptor/Met type protein kinase, is a cell-surface proto-oncogene that mediates the biological activities of hepatocyte growth factor. The protein is a disulfide-linked heterodimer comprised of two subunits: a 50 kD extracellular alpha subunit and a membrane-spanning 145 kD beta subunit that binds ligand. Met is an important regulator of cell proliferation and differentiation, organ regeneration, embryogenesis, and tumorigenesis.
Product Categories/Family for anti-MET antibody	Family: Protein Kinase; Subfamily: HGF Receptor/MET

NCBI AND UNIPROT INFORMATION

NCBI GI #	42741655
NCBI GeneID	4233
NCBI Accession #	NP_000236.2
NCBI GenBank Nucleotide #	NM_000245.2
UniProt Accession #	P08581; O60366; Q12875; Q9UDX7; Q9UPL8; A1L467; B5A932; E7EQ94
Molecular Weight	85,745 Da
NCBI Official Full Name	hepatocyte growth factor receptor isoform b
NCBI Official Synonym Full Names	MET proto-oncogene, receptor tyrosine kinase
NCBI Official Symbol	MET
NCBI Official Synonym Symbols	HGFR; AUTS9; RCCP2; c-Met
NCBI Protein Information	hepatocyte growth factor receptor; SF receptor; HGF receptor; HGF/SF receptor; proto-oncogene c-Met; scatter factor receptor; tyrosine-protein kinase Met; met proto-oncogene tyrosine kinase
UniProt Protein Name	Hepatocyte growth factor receptor
UniProt Gene Name	MET
UniProt Synonym Gene Names	HGF receptor; SF receptor
UniProt Entry Name	MET_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/c-met-antibody-immunohistochemistry-elisa/51007>