

Goat anti-Human CD14 Polyclonal Antibody | anti-CD14 antibody

Goat Polyclonal to Human CD14

CATALOG # AAA51011	HOST Goat	REACTIVITY Human
------------------------------	---------------------	----------------------------

PRODUCT SYNONYMS



CD14; Polyclonal Antibody; Goat Polyclonal to Human CD14; Anti-CD14 Antibody (Internal) IHC-plus; CD14 molecule; CD14 antigen; Human CD14; anti-CD14 antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Human
Clonality	Polyclonal
Specificity	Human CD14
Purity/Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Form/Format	TBS, pH 7.3, 0.02% Sodium Azide, 0.5% BSA
Concentration	0.5mg/ml (varies by lot)
Target	CD14
Immunogen Species	CD14 antibody was raised against Human
Immunogen	CD14 antibody was raised against peptide with sequence C-KRVDADADPRQYAD, from the internal region of the protein sequence according to NP_000582.1.
Epitope	aa71-84
Disclaimer	Due to the highly specific nature of antibodies and antigens, we cannot predict or be held responsible with respect to how this antibody will behave in your systems. Researchers using this antibody should conduct optimization studies to achieve the most optimal result possible for their intended application.

Recommended Immunohistochemistry Protocol	The following protocol is a recommendation only, and AAA Biotech makes no guarantee of the results: Tissue Preparation: Formalin fixation and embedding in paraffin wax. Tissue Sectioning: Make 4-um sections and place on pre-cleaned and charged microscope slides. Heat in a tissue-dryingoven for 45 minutes at 60°C. Deparaffinization: Wash dry slides in 3 changes of xylene - 5 minutes each @ RT Rehydration: Wash slides in 3 changes of 100% alcohol - 3 minutes each @ RT Wash slides in 2 changes of 95% alcohol - 3 minutes each @ RT Wash slides in 1 change of 80% alcohol - 3 minutes @ RT Rinse slides in gentle running distilled water - 5 minutes @ RT Antigen retrieval: Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes Remove from heat and let stand at room temperature in buffer - 20 minutes Rinse in 1X TBS with Tween (TBST) -1 minute @ RT Immunostaining: (Do not allow tissues to dry at any time during the staining procedure) Apply a universal protein block - 20 minutes @ RT Drain protein block from slides, apply diluted primary antibody - 45 minutes @ RT Rinse slides in 1 X TBST - 1 minute @ RT Apply a biotinylated secondary antibody appropriate for the primary antibody - 30 minutes @ RT Rinse slides in 1X TBST -1 minute @ RT Apply alkaline phosphatase streptavidin - 30 minutes @ RT Rinse slides in 1X TBST -1 minute @ RT Apply alkaline phosphatase chromogen substrate - 30 minutes @ RT Wash slides in distilled water - 1 minute @ RT Dehydrate: (This method should only be used if the chromogen substrate is alcohol insoluble (e.g. Vector Red, DAB) Wash slides in 2 changes of 80% alcohol - 1 minute each @ RT Wash slides in 2 changes of 95% alcohol - 1 minute each @ RT Wash slides in 3 changes of 100% alcohol - 1 minute each @ RT Wash slides in 3 changes of xylene - 1 minute each @ RT Apply coverslip
Preparation and Storage	Aliquot and store at -20°C. Avoid freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)
Anti-CD14 antibody IHC of human spleen. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 4 ug/ml.

WB (Western Blot)
1 ug/ml) Staining of Human Lymph Node lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

ADDITIONAL INFORMATION

Related Product Information for anti-CD14 antibody	CD14 is a surface protein preferentially expressed on monocytes/macrophages. It binds lipopolysaccharide binding protein and recently has been shown to bind apoptotic cells.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557417
NCBI GeneID	929
NCBI Accession #	NP_000582.1
NCBI GenBank Nucleotide #	NM_000591.3
UniProt Accession #	P08571; Q53XT5; Q96FR6; Q96L99; Q9UNS3
NCBI Official Full Name	monocyte differentiation antigen CD14
NCBI Official Synonym Full Names	CD14 molecule
NCBI Official Symbol	CD14
NCBI Protein Information	monocyte differentiation antigen CD14; myeloid cell-specific leucine-rich glycoprotein
UniProt Protein Name	Monocyte differentiation antigen CD14
UniProt Gene Name	CD14
UniProt Entry Name	CD14_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cd14-antibody-elisa/51011>