

Mouse Cystatin C ELISA Kit | CST3 elisa kit

Cystatin C Mouse ELISA Kit

CATALOG # AAA58886	REACTIVITY Mouse	APPLICATIONS ELISA
------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

Cystatin C; Cystatin C Mouse ELISA Kit; Mouse Cystatin C; CST3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Assay Type	Sandwich
Detection Range	312 pg/ml-20,000 pg/ml
Sensitivity	< 10 pg/ml
Applicable Applications for CST3 elisa kit	ELISA
Preparation and Storage	Store at 2-8 degree C for 4 months, or at -20 degree C for 8 months.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CST3 elisa kit	Principle of the Assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-Cystatin C polyclonal antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-Cystatin C polyclonal antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the Cystatin C amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of Cystatin C can be calculated. Background: Cystatin C or cystatin 3, a protein encoded by the CST3 gene, is mainly used as a biomarker of kidney function. It is a single 120-residue polypeptide with a molecular mass of approximately 13.26 kD. Cystatin C plays a role in network reorganization in the epileptic dentate gyrus, especially in granule cell dispersion and guidance of migrating newborn granule cells. It is an alternative measure of kidney function that may have prognostic importance among elderly persons who do not meet standard criteria for chronic kidney disease. Among elderly persons without chronic kidney disease, cystatin C is a prognostic biomarker of risk for death, cardiovascular disease, and chronic kidney disease, thereby signaling a state of 'preclinical' kidney dysfunction.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	755738
NCBI GeneID	1471
UniProt Accession #	P01034; Q6FGW9; B2R5J9; D3DW42
Molecular Weight	15,799 Da
NCBI Official Full Name	cystatin C
NCBI Official Synonym Full Names	cystatin C
NCBI Official Symbol	CST3
NCBI Official Synonym Symbols	ARMD11
NCBI Protein Information	cystatin-C; cystatin 3; cystatin-3; gamma-trace; post-gamma-globulin; bA218C14.4 (cystatin C); neuroendocrine basic polypeptide
UniProt Protein Name	Cystatin-C
UniProt Gene Name	CST3
UniProt Entry Name	CYTC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.