

Human ICAM-1 / CD54 ELISA Kit | ICAM-1 / CD54 elisa kit

ICAM-1 / CD54 Human ELISA Kit

CATALOG # AAA58915	REACTIVITY Human	APPLICATIONS ELISA
------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

ICAM-1 / CD54; ICAM-1 / CD54 Human ELISA Kit; Human ICAM-1 / CD54; ICAM-1 / CD54 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Assay Type	Sandwich
Detection Range	156 pg/ml-10,000 pg/ml
Sensitivity	< 10pg/ml
Applicable Applications for ICAM-1 / CD54 elisa kit	ELISA
Preparation and Storage	Store at 2-8 degree C for 4 months, or at -20 degree C for 8 months.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ICAM-1 / CD54 elisa kit

Principle of the assay: This kit was based on sandwich enzyme-linked immune-sorbent assay technology. Anti-ICAM-1 polyclonal antibody was pre-coated onto 96-well plates. And the biotin conjugated anti-ICAM-1 polyclonal antibody was used as detection antibodies. The standards, test samples and biotin conjugated detection antibody were added to the wells subsequently, and wash with wash buffer. Avidin-Biotin-Peroxidase Complex was added and unbound conjugates were washed away with wash buffer. TMB substrates were used to visualize HRP enzymatic reaction. TMB was catalyzed by HRP to produce a blue color product that changed into yellow after adding acidic stop solution. The density of yellow is proportional to the ICAM-1 amount of sample captured in plate. Read the O.D. absorbance at 450nm in a microplate reader, and then the concentration of ICAM-1 can be calculated.

Background: ICAM-1 (Intercellular Adhesion Molecule 1) also known as CD54 (Cluster of Differentiation 54) is a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by rhinovirus as a receptor. ICAM-1 is a member of the immunoglobulin superfamily, it can be induced by interleukin-1 (IL-1) and tumor necrosis factor alpha (TNFalpha) and is expressed by the vascular endothelium, macrophages, and lymphocytes. It is also an endothelial- and leukocyte-associated transmembrane protein long known for its importance in stabilizing cell-cell interactions and facilitating leukocyte endothelial transmigration. More recently, ICAM-1 has been characterized as a site for the cellular entry of human rhinovirus.

NCBI AND UNIPROT INFORMATION

NCBI GI #	167466198
NCBI GeneID	3383
NCBI Accession #	NP_000192.2
NCBI GenBank Nucleotide #	NM_000201.2
UniProt Accession #	P05362; Q5NKV7; Q96B50; B2R6M3
Molecular Weight	57,825 Da
NCBI Official Full Name	intercellular adhesion molecule 1
NCBI Official Synonym Full Names	intercellular adhesion molecule 1
NCBI Official Symbol	ICAM1
NCBI Official Synonym Symbols	BB2; CD54; P3.58
NCBI Protein Information	intercellular adhesion molecule 1; ICAM-1; cell surface glycoprotein P3.58; major group rhinovirus receptor; intercellular adhesion molecule 1 (CD54), human rhinovirus receptor
UniProt Protein Name	Intercellular adhesion molecule 1
UniProt Gene Name	ICAM1
UniProt Synonym Gene Names	ICAM-1
UniProt Entry Name	ICAM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.