

Rabbit anti-Human CD33 Monoclonal Antibody | anti-CD33 antibody

CD33 Rmab

CATALOG # AAA59308	HOST Rabbit	REACTIVITY Human
------------------------------	-----------------------	----------------------------

PRODUCT SYNONYMS

CD33; Monoclonal Antibody; CD33 Rmab; anti-CD33 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Clone Number	RBT-CD33
Sequence Length	183
Intended Use	This antibody is intended for use in Immunohistochemical applications on formalin-fixed paraffin-embedded tissues (FFPE), frozen tissue sections and cell preparations. Interpretation of results should be performed by a qualified medical professional.
Immunogen	Recombinant protein encoding the extracellular domain of human CD33.
Summary and Explanation	CD33 (gp62 or siglec-3) is a glycosylated transmembrane protein that is a member of the sialic acid-binding immunoglobulin-like lectin (siglec) family. The genomic locus of this protein has been mapped to chromosome 19q13.1-3.5. The function of CD33 is not known, but it may have a role in cell-to-cell adhesion. In maturing granulocytic cells, there is progressive down-regulation of CD33 from the blast stage to mature neutrophils. However, in monocytes and macrophages/histiocytes, strong expression of CD33 is maintained throughout maturation. Detection of CD33 using monoclonal antibodies has been a critical component in immunophenotyping acute leukemias, particularly Acute Myeloid Leukemias.
Localization	Membranous
Control	Placenta, Myometrium, Lung, Colon, Spleen, Lymph Node, Tonsil, Acute Myeloid Leukemia
Presentation	CD33 is a rabbit monoclonal antibody derived from cell culture supernatant that is concentrated, dialyzed, filter sterilized and diluted in buffer pH 7.5, containing BSA and sodium azide as a preservative.
Precautions	1. This product contains sodium azide (NaN₃), a toxic chemical which may react with plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent sodium azide build-up. 2. Ensure proper handling procedures are used with reagent. Always wear proper laboratory equipment such as laboratory coat and gloves when handling reagents. 3. Unused solution should be disposed of according to local and federal regulations. 4. Do not ingest reagent. If reagent ingested, contact a poison control center immediately. 5. For complete recommendations for handling biological specimens please refer to the CDC document, "Guidelines for Safe Work Practices in Human and Animal Laboratories" (5).

Specimen Preparation	Paraffin sections: The antibody can be used on formalin-fixed paraffin-embedded (FFPE) tissue sections. Ensure tissue undergoes appropriate fixation to ensure best results. Pre-treatment of tissues with heat-induced epitope retrieval (HIER) is recommended using ImmunoDNA Retriever with Citrate (Please Inquire), ImmunoDNA Retriever with EDTA (Please Inquire) or ImmunoDNA Digestor (Please Inquire). See reverse side for complete protocol. Tissue should remain hydrated via use Immuno/DNA Washer solutions (Please Inquire). Frozen sections and cell preparations: The antibody can be used for labeling acetone-fixed frozen sections and acetone-fixed cell preparations.
Staining Procedure	1. Cut and mount 3-5 micron formalin-fixed paraffin-embedded tissues on positive charged slides such as Hydrophilic Plus Slides (Please Inquire). 2. Air dry for 2 hours at 58° C. 3. Deparaffinize, dehydrate and rehydrate tissues. 4. Subject tissues to heat epitope retrieval using a suitable retrieval solution such as ImmunoDNA Retriever with Citrate (Please Inquire) or EDTA (Please Inquire). 5. Any of three heating methods may be used: a. TintoRetriever Pressure Cooker or Equivalent Place tissues/slides in a staining dish or coplin jar containing the ImmunoDNA Retriever with Citrate or EDTA, and place in the pressure cooker. Add 1-2 inches of distilled water to the pressure cooker and turn heat to high. Incubate for 15 minutes. Open and immediately transfer slides to room temperature. b. TintoRetriever PT Module or Water Bath Method Place tissues/slides in a pre-warmed staining dish or coplin jar containing the ImmunoDNA Retriever with Citrate or EDTA at 95°-99° C. Incubate for 30-60 minutes. c. Conventional Steamer Method Place tissues/slides in a pre-warmed staining dish or coplin jar containing the ImmunoDNA Retriever with Citrate or EDTA in a Steamer, cover and steam for 30-60 minutes. 6. After heat treatment, transfer slides in ImmunoDNA Retriever with Citrate or EDTA to room temperature and let stand for 15-20 minutes. 7. For manual staining, perform antibody incubation at ambient temperature. For automated staining methods, perform antibody incubation according to instrument manufacturer's instructions. 8. Wash slides with IHC wash buffer or DI water. 9. Continue IHC staining protocol.
Preparation and Storage	Store at 2-8 °C. Do not use after expiration date listed on package label. Temperature fluctuations should be avoided. Store appropriately when not in use, and avoid prolonged exposure to room temperature conditions.

IMAGES

Recommended IHC Protocol			
Step	ImmunoDetector AP/HRP	PolyDetector AP/HRP	PolyDetector Plus HRP
Peroxidase/AP Blocker	5 min.	5 min.	5 min
Primary Antibody	30-60 min.	30-60 min.	30-60 min.
1st Step Detection	10 min.	30-45 min.	15 min.
2nd Step Detection	10 min.	Not Applicable	15 min.
Substrate-Chromogen	5-10 min.	5-10 min.	5-10 min.
Counterstain	Varies	Varies	Varies



IHC (Immunohistochemistry)
Inset: IHC of CD33 on a FFPE Colon Tissue

NCBI AND UNIPROT INFORMATION

NCBI GI #	1032656012
NCBI GeneID	945
Molecular Weight	25,293 Da
NCBI Official Full Name	CD33 molecule transcript variant
NCBI Official Synonym Full Names	CD33 molecule

NCBI Official Symbol	CD33
NCBI Official Synonym Symbols	p67; SIGLEC3; SIGLEC-3
NCBI Protein Information	myeloid cell surface antigen CD33
UniProt Protein Name	Myeloid cell surface antigen CD33
UniProt Gene Name	CD33
UniProt Synonym Gene Names	SIGLEC3; Siglec-3
UniProt Entry Name	CD33_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/monoclonal/cd33-antibody-clone-rbt-cd33-elisa/59308>