

IgG secondary antibody

Anti-Goat IgG, DyLight488

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA63210	Chicken	Goat	Immunofluorescence, Western Blot


PRODUCT SYNONYMS

IgG; Polyclonal Secondary Antibody; Anti-Goat IgG; DyLight488; Immunoglobulin Y antibody; IgG secondary antibody

PRODUCT SPECIFICATIONS

Host	Chicken
Reactivity	Goat
Clonality	Polyclonal
Isotype	IgY
Specificity	Using serum samples detects goat IgG by Western blot. Reduced cross-reactivity to other species.
Purity/Purification	Epitope affinity purified
Form/Format	Polyclonal antibody supplied as a 200ul (2.5mg/ml) aliquot in PBS, 20% glycerol and 0.05% sodium azide.
Concentration	2mg/ml (varies by lot)
Applicable Applications for IgG secondary antibody	IF (Immunofluorescence), WB (Western Blot)
Conjugation	DyLight488
Immunogen	Full goat IgG affinity purified from domestic goat (<i>Capra hircus</i>) serum using protein G. Purity >95% by SDS-PAGE.
Preparation and Storage	Store at -20 C for long-term storage. Store at 2-8 C for up to one month.

IMAGES



Sample	WB	IHC (F)	IHC (P)	IF	ELISA	IEM
Goat	+++	ND	ND	+++	ND	ND

Reactivity Data

IF (Immunofluorescence)

Immunofluorescence in *Drosophila larva* muscle 6/7 expressing mCherry in neurons (DyGlut-mCherry) using 1st AbAnti-mCherry at 1/1,000 and 2nd AbAnti-goat IgY conjugated to DyLight488 at 1/500;

ADDITIONAL INFORMATION

Related Product Information for IgG secondary antibody	Chicken polyclonal antibody to goat IgG conjugated to DyLight488.
--	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/igg-secondary-antibody-immunofluorescence-western-blot-elisa/63210>