

Rat Epiregulin (EREG) ELISA Kit | EREG elisa kit

Rat Epiregulin (EREG) ELISA Kit

CATALOG #
AAA63337

REACTIVITY
Rat


PRODUCT SYNONYMS

Epiregulin (EREG); Rat Epiregulin (EREG) ELISA Kit; EPR; ER; Proepiregulin; EREG elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of EREG. No significant cross-reactivity or interference between EREG and analogues was observed.
Sequence Length	162
Samples	Serum, plasma, tissue homogenates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	6.1pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level EREG were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level EREG were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for EREG elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of EREG in rat serum, plasma, tissue homogenates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to EREG. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to EREG. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain EREG, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of EREG in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	11078689
NCBI GeneID	59325
NCBI Accession #	NP_067721.1
NCBI GenBank Nucleotide #	NM_021689.1
UniProt Accession #	Q9Z0L5
Molecular Weight	18,404 Da
NCBI Official Full Name	proepiregulin preproprotein
NCBI Official Synonym Full Names	epiregulin
NCBI Official Symbol	Ereg
NCBI Protein Information	proepiregulin
UniProt Protein Name	Proepiregulin
UniProt Gene Name	Ereg
UniProt Synonym Gene Names	EPR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.