

Mouse Protein Kinase C Epsilon (PKCe) ELISA Kit | PKCe elisa kit

Mouse Protein Kinase C Epsilon (PKCe) ELISA Kit

CATALOG #
AAA63986

REACTIVITY
Mouse


PRODUCT SYNONYMS

Protein Kinase C Epsilon (PKCe); Mouse Protein Kinase C Epsilon (PKCe) ELISA Kit; ANXV; ANX5; ANX-V; ENX2; PP4; ANX-A5; Annexin V; PKCe elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of PKCe. No significant cross-reactivity or interference between PKCe and analogues was observed.
Sequence Length	737
Samples	Tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.116ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level PKCe were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level PKCe were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PKCe elisa kit	Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of PKCe in mouse tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to PKCe. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to PKCe. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PKCe, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of PKCe in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	6755084
NCBI GeneID	18754
NCBI Accession #	NP_035234.1
NCBI GenBank Nucleotide #	NM_011104.3
UniProt Accession #	P16054
Molecular Weight	83,561 Da
NCBI Official Full Name	protein kinase C epsilon type
NCBI Official Synonym Full Names	protein kinase C, epsilon
NCBI Official Symbol	Prkce
NCBI Official Synonym Symbols	Pkce; PKC[e]; R75156; PKCepsilon; 5830406C15Rik
NCBI Protein Information	protein kinase C epsilon type
UniProt Protein Name	Protein kinase C epsilon type
UniProt Gene Name	Prkce
UniProt Synonym Gene Names	Pkce; Pkcea

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-elisa-kits/protein-kinase-c-epsilon-pkce/63986>