

Goat Tumor Necrosis Factor Alpha (TNFa) ELISA Kit | TNFa elisa kit

Goat Tumor Necrosis Factor Alpha (TNFa) ELISA Kit

CATALOG #
AAA64555

REACTIVITY
Goat


PRODUCT SYNONYMS

Tumor Necrosis Factor Alpha (TNFa); Goat Tumor Necrosis Factor Alpha (TNFa) ELISA Kit; DIF; TNF-A; TNFSF2; Cachectin; Tumor Necrosis Factor Ligand Superfamily Member 2; TNFa elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Goat
Specificity	This assay has high sensitivity and excellent specificity for detection of TNFa. No significant cross-reactivity or interference between TNFa and analogues was observed.
Sequence Length	234
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	5.9pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TNFa were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TNFa were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TNFa elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of TNFa in goat serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to TNFa. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to TNFa. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TNFa, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of TNFa in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	556695474
NCBI GeneID	100861232
NCBI Accession #	NP_001273371.1
NCBI GenBank Nucleotide #	NM_001286442.1
UniProt Accession #	P13296; Q28320; Q9MYZ2
Molecular Weight	25,519 Da
NCBI Official Full Name	tumor necrosis factor
NCBI Official Symbol	TNF
NCBI Official Synonym Symbols	TNFA; TNF-a; TNFSF2; Cachectin; TNF-alpha
NCBI Protein Information	tumor necrosis factor
UniProt Protein Name	Tumor necrosis factor
UniProt Gene Name	TNF
UniProt Synonym Gene Names	TNFA; TNFSF2; TNF-a; NTF; ICD1; ICD2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/goat-elisa-kits/tumor-necrosis-factor-alpha-tnfa/64555>