

Chicken Brain Derived Neurotrophic Factor (BDNF) ELISA Kit | BDNF elisa kit

Chicken Brain Derived Neurotrophic Factor (BDNF) ELISA Kit

CATALOG #
AAA64847

REACTIVITY
Chicken


PRODUCT SYNONYMS

Brain Derived Neurotrophic Factor (BDNF); Chicken Brain Derived Neurotrophic Factor (BDNF) ELISA Kit; Neurotrophin; Abreineurin; BDNF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Chicken
Specificity	This assay has high sensitivity and excellent specificity for detection of BDNF. No significant cross-reactivity or interference between BDNF and analogues was observed.
Sequence Length	246
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.065ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level BDNF were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level BDNF were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for BDNF elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of BDNF in chicken serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to BDNF. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to BDNF. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain BDNF, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of BDNF in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	71979935
NCBI GeneID	396186
NCBI Accession #	NP_001026787.1
NCBI GenBank Nucleotide #	NM_001031616.1
UniProt Accession #	P25429; Q45N27
Molecular Weight	27,715 Da
NCBI Official Full Name	brain-derived neurotrophic factor
NCBI Official Synonym Full Names	brain derived neurotrophic factor
NCBI Official Symbol	BDNF
NCBI Protein Information	brain-derived neurotrophic factor
UniProt Protein Name	Brain-derived neurotrophic factor
UniProt Gene Name	BDNF
UniProt Synonym Gene Names	BDNF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/chicken-elisa-kits/brain-derived-neurotrophic-factor-bdnf/64847>