

Human Protocadherin Gamma A2 (PCDHgA2) ELISA Kit | PCDHgA2 elisa kit

Human Protocadherin Gamma A2 (PCDHgA2) ELISA Kit

CATALOG #
AAA65000

REACTIVITY
Human


PRODUCT SYNONYMS

Protocadherin Gamma A2 (PCDHgA2); Human Protocadherin Gamma A2 (PCDHgA2) ELISA Kit; PCDHgA2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of PCDHGA2. No significant cross-reactivity or interference between PCDHGA2 and analogues was observed.
Sequence Length	932
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.063ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level PCDHGA2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level PCDHGA2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PCDHga2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of PCDHGA2 in human tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to PCDHGA2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to PCDHGA2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PCDHGA2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of PCDHGA2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	11056030
NCBI GeneID	56113
NCBI Accession #	NP_061738.1
NCBI GenBank Nucleotide #	NM_018915.3
UniProt Accession #	Q9Y5H1; Q52LL6; Q9Y5D5
Molecular Weight	90,284 Da
NCBI Official Full Name	protocadherin gamma-A2 isoform 1
NCBI Official Synonym Full Names	protocadherin gamma subfamily A, 2
NCBI Official Symbol	PCDHGA2
NCBI Official Synonym Symbols	PCDH-GAMMA-A2
NCBI Protein Information	protocadherin gamma-A2
UniProt Protein Name	Protocadherin gamma-A2
UniProt Gene Name	PCDHGA2
UniProt Synonym Gene Names	PCDH-gamma-A2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.