

Human Protoporphyrinogen Oxidase (PPOX) ELISA Kit | PPOX elisa kit

Human Protoporphyrinogen Oxidase (PPOX) ELISA Kit

CATALOG #
AAA65098

REACTIVITY
Human


PRODUCT SYNONYMS

Protoporphyrinogen Oxidase (PPOX); Human Protoporphyrinogen Oxidase (PPOX) ELISA Kit; VP; PPO; V290M; Variegate Porphyria; PPOX elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of PPOX. No significant cross-reactivity or interference between PPOX and analogues was observed.
Sequence Length	477
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.053ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level PPOX were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level PPOX were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PPOX elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of PPOX in human tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to PPOX. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to PPOX. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PPOX, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of PPOX in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506001
NCBI GeneID	5498
NCBI Accession #	NP_000300.1
NCBI GenBank Nucleotide #	NM_000309.3
UniProt Accession #	P50336; Q5VTW8; D3DVG0
Molecular Weight	50,765 Da
NCBI Official Full Name	protoporphyrinogen oxidase
NCBI Official Synonym Full Names	protoporphyrinogen oxidase
NCBI Official Symbol	PPOX
NCBI Official Synonym Symbols	VP; PPO; V290M
NCBI Protein Information	protoporphyrinogen oxidase
UniProt Protein Name	Protoporphyrinogen oxidase
UniProt Gene Name	PPOX
UniProt Synonym Gene Names	PPO

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.