

Mouse Syndecan 1 (SDC1) ELISA Kit | SDC1 elisa kit

Mouse Syndecan 1 (SDC1) ELISA Kit

CATALOG # AAA65686	REACTIVITY Mouse
------------------------------	----------------------------


PRODUCT SYNONYMS

Syndecan 1 (SDC1); Mouse Syndecan 1 (SDC1) ELISA Kit; CD138; SDC; SYND1; Syndecan Proteoglycan 1; SDC1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of SDC1. No significant cross-reactivity or interference between SDC1 and analogues was observed.
Sequence Length	311
Assay Type	Quantitative Sandwich
Samples	Mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Detection Range	125-8000pg/mL
Sensitivity	< 47pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SDC1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SDC1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SDC1 elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SDC1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SDC1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SDC1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of SDC1 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of SDC1 in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	6755438
NCBI GeneID	20969
NCBI Accession #	NP_035649.1
NCBI GenBank Nucleotide #	NM_011519.2
UniProt Accession #	P18828; Q62278; Q9WVD2
Molecular Weight	28,235 Da
NCBI Official Full Name	syndecan-1
NCBI Official Synonym Full Names	syndecan 1
NCBI Official Symbol	Sdc1
NCBI Official Synonym Symbols	Sstn; Synd; CD138; Synd1; syn-1; AA408134; AA409076
NCBI Protein Information	syndecan-1
UniProt Protein Name	Syndecan-1
UniProt Gene Name	Sdc1
UniProt Synonym Gene Names	Synd-1; Synd1; SYND1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/mouse-elisa-kits/syndecan-1-sdc1/65686>