

Human Fibronectin (FN) ELISA Kit | FN elisa kit

Human Fibronectin (FN) ELISA Kit

CATALOG #
AAA65869

REACTIVITY
Human


PRODUCT SYNONYMS

Fibronectin (FN); Human Fibronectin (FN) ELISA Kit; HSP70-1; HSP-A1A; HSPA1; HSPA1B; HSP70-2; FN elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of FLG. No significant cross-reactivity or interference between FLG and analogues was observed.
Sequence Length	2446
Samples	Human Tissue Homogenates, Cell Lysates, Cell Culture Supernates Or Other Biological Fluids
Assay Type	Quantitative Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.114ng/mL.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FLG were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FLG were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FN elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of FLG in human tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to FLG. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to FLG. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FLG, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of FLG in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	807066342
NCBI GeneID	2335
NCBI Accession #	NP_001293058.1
NCBI GenBank Nucleotide #	NM_001306129.1
UniProt Accession #	P02751; O95609; O95610; Q14312; Q14325; Q14326; Q17RV7; Q564H7; Q585T2; B7ZLF0; E9PE77; E9PG29
Molecular Weight	256,502 Da
NCBI Official Full Name	fibronectin isoform 8
NCBI Official Synonym Full Names	fibronectin 1
NCBI Official Symbol	FN1
NCBI Official Synonym Symbols	FN; CIG; FNZ; MSF; ED-B; FINC; GFND; LETS; GFND2
NCBI Protein Information	fibronectin
UniProt Protein Name	Fibronectin
UniProt Gene Name	FN1
UniProt Synonym Gene Names	FN; FN; CIG

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.