

Mouse Carnitine Acetyltransferase (CRAT) ELISA Kit | CRAT elisa kit

Mouse Carnitine Acetyltransferase (CRAT) ELISA Kit

CATALOG #
AAA66051

REACTIVITY
Mouse


PRODUCT SYNONYMS

Carnitine Acetyltransferase (CRAT); Mouse Carnitine Acetyltransferase (CRAT) ELISA Kit; CAT1; Carnitine O-Acetyltransferase; Carnitine acetylase; CRAT elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of CRAT. No significant cross-reactivity or interference between CRAT and analogues was observed.
Sequence Length	626
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.047ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CRAT were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CRAT were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CRAT elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CRAT in mouse tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CRAT. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CRAT. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CRAT, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CRAT in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	85662408
NCBI GeneID	12908
NCBI Accession #	NP_031786.2
NCBI GenBank Nucleotide #	NM_007760.3
UniProt Accession #	P47934; Q3TCG3; Q3V1Y3; Q923A6
Molecular Weight	70,840 Da
NCBI Official Full Name	carnitine O-acetyltransferase
NCBI Official Synonym Full Names	carnitine acetyltransferase
NCBI Official Symbol	Crat
NCBI Official Synonym Symbols	CARAT; AW107812
NCBI Protein Information	carnitine O-acetyltransferase
UniProt Protein Name	Carnitine O-acetyltransferase
UniProt Gene Name	Crat
UniProt Synonym Gene Names	Carnitine acetylase; CAT; CrAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.